

---

# JMIR XR and Spatial Computing (JMXR)

---

Volume 2 (2025) ISSN 2818-3045 Editor in Chief: Lars Riedemann, MD

---

## Contents

### Review

Immersive Virtual Reality for Health Promotion and Primary Prevention in Psychology: Scoping Review  
([e49923](#))  
Solenne Bonneterre, Oulmann Zerhouni, Marilisa Boffo. . . . . 2

### Original Paper

Augmented Reality in Enhancing Operating Room Crisis Checklist Adherence: Randomized Comparative  
Efficacy Study ([e60792](#))  
Rayan Ebnali Harari, Abdullah Altaweel, Erik Anderson, Charles Pozner, Rafael Grossmann, Andrew Goldsmith, Hamid Shokoohi. . . . . 18

# Immersive Virtual Reality for Health Promotion and Primary Prevention in Psychology: Scoping Review

Solenne Bonneterre<sup>1</sup>, PhD; Oulmann Zerhouni<sup>2</sup>, Prof Dr; Marilisa Boffo<sup>3</sup>, PhD

<sup>1</sup>Laboratoire Parisien de Psychologie Sociale, Département de Psychologie, Université Paris Nanterre, Nanterre, France

<sup>2</sup>Centre de Recherche sur les Fonctionnements et Dysfonctionnements Psychologiques, UR 7475, Université de Rouen Normandie, Rouen, France

<sup>3</sup>Department of Psychology, Education, and Child Studies, Erasmus School of Social and Behavioral Sciences, Erasmus University Rotterdam, Rotterdam, Netherlands

## Corresponding Author:

Solenne Bonneterre, PhD

Laboratoire Parisien de Psychologie Sociale, Département de Psychologie, Université Paris Nanterre, Nanterre, France

## Abstract

**Background:** Virtual reality (VR) has emerged as a promising tool in health promotion and prevention psychology. Its ability to create immersive, engaging, and standardized environments offers unique opportunities for interventions and assessments. However, the scope of VR applications in this field remains unclear.

**Objective:** This scoping review aims to identify and map the applications of VR in health promotion and prevention psychology, focusing on its uses, outcomes, and challenges.

**Methods:** A systematic search was conducted across 3 electronic databases (PubMed, PsycINFO, and Scopus) for studies published between 2010 and 2024. Eligibility criteria included empirical studies using immersive VR for health promotion and prevention, while studies using nonimmersive VR, lacking health-related applications, or focusing on clinical interventions were excluded. The review followed PRISMA-ScR (Preferred Reporting Items for Systematic Reviews and Meta-analyses extension for Scoping Reviews) guidelines, and 4295 records were initially identified, with 51 studies included after screening. Data were synthesized qualitatively to identify key applications, limitations, and emerging trends.

**Results:** VR was primarily used in three areas: (1) delivering interventions (eg, pilot testing, skills training), (2) exploring fundamental research questions, and (3) assessing outcomes such as behavioral or psychological responses. Although VR demonstrated potential for enhancing user engagement and replicating ecological scenarios, its effectiveness compared to nonimmersive methods varied. Most studies were pilot or feasibility studies with small, nonrepresentative samples, short follow-up periods, and limited methodological standardization.

**Conclusions:** VR offers a versatile and promising tool for health promotion and prevention but its applications are still in the early stages. The evidence is limited by methodological weaknesses and variability in outcomes. Future research should prioritize replication, longitudinal designs, and standardized methodologies to strengthen the evidence base and expand the applicability of VR interventions.

(*JMIR XR Spatial Comput* 2025;2:e49923) doi:[10.2196/49923](https://doi.org/10.2196/49923)

## KEYWORDS

virtual reality; health psychology; prevention psychology; health promotion

## Introduction

### Background

Health and prevention psychology aims to address health-related issues to either prevent individuals from starting or continuing an unhealthy behavior (ie, primary prevention), help them to detect or reduce illness in early stages (ie, secondary prevention), or support individuals in their journey against consequences of heavier injuries or diseases (ie, tertiary prevention, [1]). Although secondary and tertiary prevention are more individual-based depending on the illness or signs or symptoms individuals need to learn to cope with, primary prevention is

broader and aimed at a larger audience. Therefore, primary or universal prevention is designed to prevent individuals from the general population from getting injured or sick and aims to enable people to live a sustainable and healthy lifestyle [2,3].

In this sense, health promotion campaigns have started to integrate technological innovations such as virtual reality (VR). We refer to VR as a type of human-computer interface immersing users into a computer-generated 3D virtual environment (VE) they can interact with in a naturalistic fashion, usually via an avatar (ie, representation of the user in the VE [4]). More pragmatically, we labeled as VR any type of device that has the ability to sensorily detach the user from the outside

world (at least sight, but also sounds, smell, and touch in some cases). This includes the use of a cave automatic VE (users are surrounded by walls displaying the VE) or a head-mounted display (HMD), which blocks the user's field of view outside of the VE and from which the user cannot turn away by simply looking away (ie, computer screens or 360° videos will not be considered VR in this definition).

The main aim of VR is to recreate a realistic, ecological context and experience while keeping some degree of experimental control over it [5-7]. Systematic reviews have reported promising results from VR-based interventions in other disciplines (eg, clinical psychology [8] and social psychology [9,10]). However, to our knowledge, there has been no review of the use of VR technologies for primary health promotion and prevention. Therefore, instead of focusing on specific research questions related to a topic, outcome, or population, the goal of this review was to map the current state of the art of the use of VR in such areas and identify gaps and future directions.

## Rationale

### *Virtual Reality: Operating Principles*

The VR literature highlights 2 essential concepts, immersion and presence, both of which are critical to the user's experience in VEs [7,11]. Immersion refers to the technological ability of a VR system to fully engage the user by replacing real-world sensory inputs with virtual stimuli. The more immersive the device, the less interface there is between the user and the virtual world. High immersion includes naturalistic interactions, such as the use of body suits to track movement, which increases the sense of realism [7]. Immersive systems create a sense that the virtual world is an actual experience rather than a mediated one. However, presence depends on the user's psychological response to the VE. It is the subjective feeling of "being there" in the virtual world, interacting with it as if it were real [12]. This sense of presence increases engagement and leads to more vivid, memorable experiences [13]. Notably, presence can be felt in both immersive and nonimmersive media, such as movies or books, as it is influenced by individual factors and not just the technological features of the medium [14].

Although immersion and presence are often related, they are not the same. Higher levels of immersion tend to enhance feelings of presence, but immersion is not a necessary condition for presence [15]. Thus, immersion can be viewed as a moderator that enhances presence but does not guarantee it [16].

### **Why Use VR in Health Promotion and Prevention Psychology?**

VR technology has emerged as a promising tool in health promotion and prevention psychology, allowing for immersive experiences that can enhance user engagement and motivation [17,18]. VR enables researchers to create safe, ecological, and standardized VEs, where health promotion interventions can be effectively delivered and evaluated. VR presents key advantages as a tool for research and intervention in health promotion and primary prevention [7].

First, VR can be combined with devices aimed at mimicking more natural movements (eg, the use of handheld controllers

or haptic devices instead of a mouse and keyboard) and can encompass the integration of full-body motor and haptic feedback when using a bodysuit. This freedom and wholeness of movement can help enhance learning through direct practice, visualization, and ultimately embodied cognition (ie, cognition linked to the body [7,19]). Hence, VR can be a relevant tool to create interventions aimed at learning health-related behaviors that require practicing skills (eg, detecting testicular disorders [20]).

Second, due to its ability to elicit embodiment, VR is well suited to elicit and enhance perspective-taking and empathy [7,21]. For example, embodying an obese avatar could enhance taking the perspective of being overweight, leading to a more effective learning of the consequences of obesity and, in turn, a greater intention to take care of individual health (ie, reduce the attitude-intention-behavior gap). Through the feeling of presence, individuals can visualize themselves in a specific situation, hence allowing a deeper sense of self-reflection [20], potentially leading to more persistent changes in behavior. VR can recreate ecological situations and environments in which users can embody an avatar and act in the virtual world as if it were real, through the feeling of presence [12].

## Objective

Our goal was to identify and map how VR has been used in the field of health promotion and primary prevention. In this scoping review, we addressed three broad research questions:

1. What are the uses of VR technology in primary prevention and health promotion (ie, an overview of the goals and research questions addressed through the use of VR)?
2. What do we know so far about the effects of using VR in these fields (ie, a summary of the results)?
3. What are the challenges and limitations, if any, encountered so far?

Based on the findings of the scoping review, we drafted a list of recommendations and perspectives for the use of VR in health promotion and primary prevention.

## Methods

### **Protocol and Registration**

The scoping review protocol was drafted according to the PRISMA-ScR (Preferred Reporting Items for Systematic Reviews and Meta-analyses extension for Scoping Reviews) checklist [22,23]. We also conducted a synthesis without meta-analysis [24] (Checklist 1).

### **Eligibility Criteria**

We included any peer-reviewed and published empirical article, written in English, that described a study conducted on human subjects deploying any kind of immersive VR device (eg, HMD, cave automatic VE), including 360° videos when used in a VR setup, focusing on any research question in the field of health promotion or primary prevention, from January 1, 2010, to September 16, 2024. We chose to limit the search to the last 14 years in order to generate a recent state-of-the-art overview of the field. We excluded studies conducted on nonhumans or focused on secondary or tertiary prevention interventions, such

as psychotherapeutic treatments (eg, VR exposure therapy) and medical interventions (eg, rehabilitation), or specialized educational programs unrelated to prevention (eg, skills improvement for health practitioners). Pilot studies were not excluded from this review because of their critical role in assessing the feasibility and acceptability of interventions that may inform future primary, secondary, or tertiary prevention efforts. We excluded studies using the term “virtual reality” that described computer-based VEs involving a virtual world (eg, Second Life) or computer-related or motion-sensing devices (eg, Kinect, joystick) when they were associated with a nonimmersive VR setup (eg, non-VR video or serious game). We also used the population-concept-context framework to define our inclusion criteria. The population includes adolescents, young adults, and specific populations at risk for health issues (eg, individuals with anxiety or those at risk for substance use). The concept focuses on the application of VR technology to promote health behaviors, enhance knowledge, and improve emotional well-being. The context refers to contextual factors including the environments where VR interventions are delivered, such as schools, community centers, or health care facilities.

### Information Sources and Search Process

We searched 3 databases from January 1, 2010, until September 16, 2024 (PubMed and PsycINFO). For each database, we combined 2 sets of keywords; the first set focused on health promotion and prevention psychology. For PubMed, the search strings were (“health prevention” OR “health promotion” OR “health risk communication” OR “health communication” OR “preventive psychology” OR “behavior change” OR “attitude change”) AND (“virtual reality” OR “immersive virtual reality” OR “immersive virtual environment”). For PsycINFO, the search strings were (“health prevention” OR “health promotion” OR “health risk communication” OR “health communication” OR “preventive psychology” OR “behavior change” OR “attitude change”) AND (“virtual reality” OR “immersive virtual reality” OR “immersive virtual environment”).

### Selection of Sources of Evidence

Studies that did not employ VR technology, were not peer-reviewed, were reviews or meta-analyses, or lacked empirical data were excluded from the review. The screening process was conducted in 2 stages to enhance the rigor of the selection. In the first stage, titles and abstracts of the identified studies were reviewed to determine their relevance based on the inclusion criteria. This initial screening allowed the authors to eliminate studies that were clearly outside the scope of the review. In the second stage, full-text articles of the remaining studies were assessed to confirm their eligibility for inclusion. The extraction process was conducted independently by multiple reviewers to enhance reliability and minimize bias. Any discrepancies in data extraction were resolved through discussion and consensus among the reviewers. This meticulous approach to data extraction allowed the authors to synthesize findings across studies effectively and draw meaningful conclusions regarding the efficacy and feasibility of VR interventions in health promotion and primary prevention.

### Data Charting Process

The data charting process involved collecting information on study characteristics, intervention details, measured outcomes, user experience, type of materials, and sample characteristics (see [Multimedia Appendix 1](#)). Of note, approximately 63% of the studies included in the review were categorized as pilot or feasibility studies. We also recorded the type of VR technology used (eg, immersive headsets, desktop VR), the duration of the intervention, and the focus of the VR content (eg, health education, behavior change). On average, participants spent approximately 12.8 (SD 11.1) minutes using VR. We focused on health-related outcomes such as knowledge acquisition, behavioral intentions, and psychological well-being. User experience was assessed through qualitative data that provided insights into participants’ enjoyment, ease of use, and perceived effectiveness of the VR interventions. Many studies found that participants found the VR experience both enjoyable and engaging, which in turn led to higher participation rates compared to non-VR interventions.

### Data Items

Primary variables included study characteristics such as authorship, year of publication, study design, and sample size, which provided context for the research findings. Participant demographics, including age, gender, and health status, were also collected to understand the populations included in the studies. Intervention details were documented, focusing on the type of VR technology used, the duration of the intervention, and the specific health issues addressed. Measured outcomes were categorized into primary outcomes, such as knowledge acquisition and behavioral intentions, and secondary outcomes, including user engagement and satisfaction. User experience data were collected to assess participants’ enjoyment, ease of use, and any challenges encountered during the VR interventions. In addition, limitations of the studies were noted, including issues such as small sample sizes and methodological limitations, which are critical for contextualizing the findings. It is important to note that while immersion and presence are key concepts in understanding the effectiveness of VR, these variables were not measured consistently across studies, which may affect the interpretation of results. The data elements collected were intended to provide a structured review of the existing literature, as well as identify trends, gaps, and implications for future research in the field of VR-based health interventions.

### Critical Appraisal of Individual Sources of Evidence

We found that approximately 37% of the included studies were pilot or feasibility studies. These studies primarily focused on evaluating the usability and acceptability of VR interventions, which are critical for assessing the feasibility of larger-scale research. Although pilot studies provide valuable insights into user experiences and preliminary results, their small sample sizes and limited generalizability limit the ability to draw firm conclusions about the effectiveness of VR-based interventions. Mixed results have been found when comparing VR interventions to traditional methods, suggesting that VR does not always offer a clear advantage in achieving health outcomes. Key variables such as immersion and presence, which are critical

to understanding how VR might influence health behaviors, have not been systematically evaluated. We found a lack of focus on larger, more diverse samples and aim to replicate existing studies to strengthen the evidence supporting the use of VR in health promotion efforts.

### Study Selection Procedure

All search results were stored in Zotero, an open-source reference manager, and duplicates were removed. Titles and abstracts were screened first, removing articles that clearly did not match eligibility criteria. Second, full texts of the remaining articles were downloaded to define final eligibility for inclusion. For each step, 2 reviewers conducted the screening independently and compared and discussed these discrepancies until a full consensus was reached.

### Data Extraction Process and Synthesis of Results

Data extraction was done by 1 reviewer, who extracted the following items from the included articles: (1) title and authors, (2) goal(s) of the study, (3) design of the study, (4) study sample characteristics, (5) VR device used, (6) main results, and (7) limitations reported by the authors. A second reviewer verified that all data were correctly extracted. Following the data extraction, we conducted a narrative analysis and synthesis of the results. Results and implications of the data extracted from

the included studies were discussed by 2 reviewers in relation to the 3 research questions of the scoping review.


## Results

### Study Selection and Characteristics of Included Studies


The initial search identified 4295 unique articles, which were reduced to 51 eligible articles (see the PRISMA flowchart in [Figure 1](#)). Included studies were conducted in Asia (11 studies, 22%), Europe (18 studies, 35%), the Middle East (1 study, 2%), and North America (21 studies, 41%). The total sample size across all studies was 4647 participants, with an average of 91.1 participants per study. Study samples included slightly more women, with 2651 women (53%) and 1958 men (42.7%). The mean age of participants across the studies was 31.6 (SD 5.45) years. Studies primarily included adults, with 29 studies (57%) focused on adults, followed by 14 studies (28%) focused on adolescents, 7 studies (14%) focused on senior adults, and 1 study (2%) focused on children. Specific populations studied included students (7 studies, 29%), people with cognitive impairment (3 studies, 12%), and people with obesity (3 studies, 12%). Other populations studied included former smokers (1 study, 4%), NHS staff (1 study, 4%), parents (2 studies, 8%), smokers (2 studies, 8%), adults who had been in lockdown (1 study, 4%), and unvaccinated adults (1 study, 4%, see [Figure 2](#) for details).



**Figure 1.** Flowchart of the study selection process following PRISMA guidelines. A total of 4295 articles were initially identified across 3 databases. After removing duplicates and applying eligibility criteria, 51 studies were included. iVR: immersive virtual reality; PRISMA: Preferred Reporting Items for Systematic Reviews and Meta-analyses.



**Figure 2.** Overview of key outcomes from the intervention studies.



**Characteristics of Sources of Evidence**

The 51 included studies focused on various health-related topics (Table 1), the most predominant ones being nutrition (17%) and risky behaviors (4%). All studies used HMD, except for Lemieux et al [25], where the device used was not mentioned. HMDs were mainly Oculus (Quest, Go, or Rift, 24%), HTC Vive (17%), or Samsung Gear VR (15%). Almost half of the

studies (43%) were coupled with 1 or 2 handheld controllers. Most studies (56%) included an active interaction with the VE by using 1 or 2 handheld controllers or the bodysuit to interact with the VE. About 49% of VR exposure lasted a maximum of 10 minutes, including 22% of studies with under 5 minutes of VR exposure. We estimated an average time of 12.8 (SD 11.1) minutes spent using VR, according to the information given in the articles.

**Table .** Characteristics of sources of evidence.

Area of study	Studies
Nutrition, including nutrition and obesity prevention (17%)	Blom et al [26]; Isgin-Atici et al [27]; Ledoux et al [28]; Marcum et al [29]; McBride et al [30]; Persky et al [31,32]; Verhulst et al [33]
Eating disorder and binge eating (5%)	Ferrer-Garcia et al [34]; Lemieux et al [25]
Sugar-sweetened drink consumption (10%)	Blom et al [26]; Ledoux et al [28]; Marcum et al [29]; McBride et al [30]
Smoking tobacco (8%)	Borrelli et al [35]; Ferrer-García et al [36]; García-Rodríguez et al [37]; Bonnetterre et al [17]
Smoking e-cigarettes (5%)	Weser et al [38,39]
Alcohol use (5%)	Guldager et al [40]; Ma [41]
Risk behavior in adolescents (4%)	Hadley et al [42,43]
Gambling (2%)	Detez et al [44]
Exposure to nature to enhance well-being/stress reduction (14%)	Alyan et al [45]; Beverly et al [46]; Brimelow et al [47,48]; Browning et al [49]; Calogiuri et al [50]
General well-being/stress reduction (10%)	Afifi et al [51]; Adhyaru et al [52]; Kim et al [53]; Riva et al [54]; Ko et al [55]; Kiper et al [56]
Using handheld controllers (7%)	Eisapour et al [57]; Fang and Huang [58]; Farič et al [59]
Using a connected bike (7%)	Bird et al [60]; Zeng et al [61,62]
Vaccination (4%)	Mottelson et al [63]; Nowak et al [64]
Medication-taking (2%)	Niki et al [65]

## Results of Individual Sources of Evidence: Detailed Results

### Main Identified Research Goals

We identified three main goals for using VR: (1) as a tool to deliver an intervention, with 35 articles focusing on either (1a) pilot testing or testing the feasibility of using VR materials or procedures or (1b) using VR to deliver an actual intervention (eg, skills learning, comparing VR vs other intervention modalities) to test its relative efficacy; (2) as a tool to address fundamental research questions, with 6 studies aimed at recreating ecological settings to address physiological and psychological changes when exposed to certain situations (eg, cravings elicitation); or (3) as an assessment tool, with 5 studies investigating food choices with a food buffet created in VR.

### Pilot Studies: Ensuring Usability and Enjoyability

Many studies included in the scoping review were pilot or feasibility studies (about 37%, Table 2) from which we distinguished two main purposes: (1) testing VR usability for future research and seeing how target outcomes are impacted and (2) assessing users' experience with VR. First, researchers

found that the use of VR in their methods was rather relevant and reached multiple target outcomes such as reducing stress using a short exposure to nature in VR [45-48,51,52,55], even though exposure durations were relatively short (3-10 minutes). The use of VR was also useful to enhance participants' physical and cognitive activity [66,67]. Finally, some studies were focused on prevention and the major advantage of VR use is its ability to involve participants directly in the preventive message, for example through gaming [68] or skill practice (eg, refusing peer pressure to vape [38]). This resulted in improved knowledge on health topics (eg, on smoking in [69]) and intentions to check for diseases (eg, [20]). It also helped to deliver information in a more traditional preventive way (eg, exposure to a preventive video in an HMD in [35] or a FestLab in [40]). Overall, pilot and feasibility studies, even if conducted on small samples, found VR to be enjoyed and accepted by participants, as well as useful and feasible, and found that it impacted target outcomes (eg, enhanced well-being, increased knowledge). These results occurred whether participants only had a one-time exposure (eg, [53,69]) or sessions over a few weeks (eg, [48]) and were found to be sustained at follow-up when measured (eg, participants reduced their tobacco intake over the month following their participation [35]).



**Table .** Summary of articles and their classification within the scoping review.

Category and study	Descriptives	iVR <sup>a</sup> details	Objective(s)	Study design	Main conclusions
<b>1a: Pilot or feasibility studies</b>					
Adhyaru and Kemp [52]	n=39; mean age 36.6 (SD 10.3) years; 82% women; health care workers	HMD <sup>b</sup> (Oculus Go); 10 minutes	Explore if exposure to nature in iVR can help health care workers de-stress at work.	Before-after exposure; within-subject	iVR reduced anxiety, anger, and heart rate, and enhanced happiness and relaxation.
Affi et al [51]	n=50 older adults with cognitive impairments and their family members	Immersive VR <sup>c</sup> system	Assess whether iVR improves quality of life and social interaction for older adults and their family members.	Feasibility study with pre-post assessments	VR improved social interaction and quality of life for both older adults and their families.
Alyan et al [45]	n=20; mean age 21.8 (SD 2.2) years; 50% women; students	HMD (HTC Vive); 5 minutes	Use iVR to reduce stress via a virtual walk in nature.	2 (environment: realistic vs dreamlike); between-subject	iVR reduced stress and enhanced mental well-being.
Beverly et al [46]	n=102; 72% women; health care workers	HMD (Oculus Go/Pico G2); 3 minutes	Explore if cinematic iVR can reduce stress in health care workers.	Before-after exposure; within-subject	iVR reduced stress, independently of previous iVR use or job type.
Bonnetterre et al [17]	n=121; mean age 19.6 years; 82.5% female; university students	Sensiks Immersive VR system	Evaluate the impact of VR on memorization, attitudes, and craving responses to anti-tobacco posters.	Randomized controlled trial	VR enhanced memorization of prevention messages.
Borelli et al [35]	n=23; mean age 49.8 (SD 13.3) years; 22% women; adult smokers	HMD (Knoxlabs V2 cardboard); 5 minutes	Examine the feasibility and impact of a smoking cessation intervention during dental cleaning.	2 (video type: smoker ready/not ready to quit) × 3 (time: pre/post/follow-up); within-subject	Feasible and accepted by both smokers and dental care providers.
<b>1b: Interventions</b>					
Ahn [5]	n=73; mean age 20.8 (SD 1.1) years; 82% women; students	HMD (NM); 2 minutes	Test efficacy of preventive messages on sugar and sweetened beverage consumption via avatar embodiment.	2 (pamphlet only vs pamphlet plus iVR) × 2 (tailoring: others vs self) × 3 (time: pre/post/follow-up); between-subject	iVR heightened intentions to limit sugar and sweetened beverage consumption; effects were present at follow-up.
Blom et al [26]	n=99; mean age 30.7 (SD 10.9) years; 60% women; general population	HMD (HTC Vive); ≥3 minutes	Study purchase behaviors in an iVR supermarket.	2 (nudge vs control) × 2 (time pressure: 3 minutes vs no pressure); between-subject	iVR revealed changes in healthy food purchases based on nudge type.
<b>2: Fundamental research</b>					
Chittaro et al [70]	n=105; mean age 21.49 (SD 2.43) years; 90.5% women; students	HMD (Sony HMZ-T1); 5 minutes	Investigate links between iVR and persuasion theory, including inducing mortality salience.	2 (environment: iVR park vs cemetery); between-subject	iVR elicited mortality salience, impacted attitudes, and induced greater physiological reactions than traditional mortality salience manipulations.
Ferrer-Garcia et al [36]	n=25; mean age 29.7 (SD 13.4) years; 32% women; smokers	HMD (5DT HMD 800); time not mentioned	Assess iVR's ability to produce cravings toward tobacco smoking.	Before-during exposure to smoking cues	iVR created cravings, correlated with presence.
<b>3: Assessment tool</b>					
Isgin-Atici et al [27]	n=73; mean age 22.2 (SD 4.1) years; 56% women; students	HMD (HTC Vive); 5 - 25 minutes	Evaluate ease of use and efficiency of a virtual cafeteria.	2 (groups: iVR novices vs experienced); between-subject	iVR was user-friendly and effective regardless of prior VR experience.

Category and study	Descriptives	iVR <sup>a</sup> details	Objective(s)	Study design	Main conclusions
Marcum et al [29]	n=221; mean age 38 (SD 5.6) years; 100% women; mothers with obesity	HMD; time not mentioned	Examine microbehaviors influencing food selection in an iVR buffet.	3 (conditions: food safety control vs behavioral risk information vs family-based risk information); between-subject	iVR enabled dynamic assessment of food choice behaviors.

<sup>a</sup>iVR: immersive virtual reality.

<sup>b</sup>HMD: head-mounted display.

<sup>c</sup>VR: virtual reality.

Second, most participants found VR enjoyable and fun [59,68] and quite easy to use [52]; some were asked to complete a short tutorial [27]. Even older adults were able to manipulate handheld controllers [57], but 1 study reported that the HMD is sometimes heavy for their neck to lift (1 participant dropped out because of this reason [52]). It is worth noting that some of these studies [20,59] involved the targeted population in co-designing the intervention in previous pilot studies, hence not only explicitly ensuring usability [57] but also enhancing users' satisfaction with the intervention. Co-designing an intervention with the targeted population and conducting a first pilot study on a small sample (eg, 12/33) can improve the level of satisfaction and usability of the intervention prototype, albeit ultimate user satisfaction can only be assessed following full-scale deployment of the intervention.

### Relative Efficacy of VR Interventions

Interventions (39% [20/51] of the studies included in the review) using VR focused on several targets such as enhancing well-being by simulating a walk in nature (while remaining seated [49,54] or walking on a treadmill [50]) or skill learning and practice on various health topics [40,42]. Some studies were interested in delivering preventive content [40,41,71], other studies used VR's ability to create standardized conditions to test theoretical frameworks (eg, nudge and time pressure on healthy food choice [26,40]), while still others used VR to embody a specific character in order to impact health outcomes [5,33,56].

The key element of most studies included in this group is that they often compared the use of VR with other modalities to deliver an intervention; for example, delivering preventive information in VR versus a 2D screen (eg, [41,54]) or without the use of specific technology (eg, live role-playing with an instructor [64], reading a pamphlet, [61]). Some studies also compared different depths of immersion [50,60,62].

When comparing the relative efficacy of VR with other modalities, mixed results were found. For example, even though participants exercising using VR experienced an attentional shift from exercising, meaning that individuals were usually distracted and entertained by the VR setting, leading them to actually enjoy physical exercise, it was not always sufficient to obtain greater physical involvement when compared to nonimmersive physical activities [25,58,60,62]. However, some studies found no difference in outcomes between the use of VR and 2D screens [54,71], and other studies even found that a virtual walk remained less efficient than a real walk in nature for mood enhancement [49]. Some studies, using VR only, also

found no impact of VR prevention interventions on target outcomes (eg, no change in physical self-perception when using VR to prevent eating disorders [39], no increased knowledge on alcohol [40]). Still, we note that VR was a great tool to induce changes in knowledge and intentions to adopt a behavior (eg, vaccination intention [61], smoking e-cigarettes [44]) and for skill practice [42].

A few recent studies [44,54] investigated the use of VR outside of the laboratory, recruiting participants who own VR devices at home. Portable VR devices have become more affordable, resulting in individuals being able to use them potentially anywhere and be autonomously engaged with VR-based interventions. Furthermore, both studies resulted in an improvement in the target outcomes (reduction of psychological distress [54], increase in vaccination [44]).

Overall, VR is impactful; it can create precise and standardized experimental situations (eg, embodying an obese or weight-gaining avatar [5,33]), and it is especially practical for skill practice and sometimes for physical activity. VR-based interventions have shown a higher degree of attendance in intervention sessions (ie, adherence) than the same intervention done without the use of VR [64]. However, when VR is only used to deliver information without leveraging its specific characteristics, such as immersivity and active use of the device (ie, interacting with the VE via a game [61]), it has often been found to have similar efficacy as more traditional ways to deliver information (eg, 2D screens).

### VR to Address Fundamental Health Research Questions: A Tool to Recreate Ecological Settings in the Lab

VR can recreate real-life situations in laboratories and has been used across different domains, such as gambling [36], tobacco cravings [28,37], and food cravings [34,70], as well as for mimicking specific situations inducing certain psychological states, such as mortality salience (eg, [29]). In all studies, exposure to specific cues (eg, food items, cemetery, individuals smoking) or situations (eg, being in a pub, gambling on a slot machine) elicited both physiological (eg, increased heart rate, arousal) and psychological (eg, self-reported craving) changes, whether individuals were actively (ie, interacting with the VE) or passively (ie, watching visual content) using the VR device, suggesting that the highly immersive characteristics of VR are effective at eliciting an emotional response.

However, only 1 study compared eliciting cravings using VR versus other types of devices [34], indicating VR is not better suited to trigger a craving response than 2D pictures. It might

be possible that this null effect was due to the passive use of VR in this specific study, as interacting with a cue in VR has been found to enhance cravings [28].

### ***VR as an Assessment Tool in Health-Related Interventions***

A total of 5 studies used VR as an assessment tool in the field of nutrition by recreating a virtual buffet displaying food [27,30-32,65], where participants' task was to collect a plate of food. Participants found the VR food buffet easy to use, independently of whether they already used a VR device in the past [27]. In this context, VR allows researchers to study precisely how many items and types of food were selected and in which quantity, enabling them to calculate the total calories contained in each plate more easily. It also helped to display to participants a standardized food buffet with diverse food items without constraints from a real food buffet (eg, expiration dates, flexibility in food types, reduced costs).

### **Study Limitations**

#### ***The Necessity to Adapt the Use of VR to Experimental Needs***

The use of VR, whether for applied or fundamental research, has shown some limitations, mainly related to the study methodology and VR technology itself (eg, cybersickness, notably in [50]). First, a majority of included studies suffered from either small sample sizes (eg, 10 participants in [65], 6 in [57]) or nonrepresentative samples (eg, students in [29], healthy and active young individuals in [58]), limiting the validity and generalizability of results. Second, the quality of the experimental designs was sometimes limited (eg, semiexperimental design with pre-post comparisons) because of a lack of a proper control condition or not conducting a rigorous randomized controlled trial [20,46]. Short-term follow-up or the lack of a follow-up altogether was also mentioned as a limiting factor in numerous studies [41,42].

#### ***Lack of Systematic Assessment of VR's Main Characteristics: Presence, Immersion, and Cybersickness***

VR's effects, especially persuasive effects, seem to come from its ability to enhance presence, which is the feeling of being there during a VR experience. Hence, participants act similarly to real life in the VE because they are fully immersed in their interaction with it. The level of presence experienced by users can impact targeted variables in the intervention; participants who felt more present in the VE showed stronger positive effects on persuasion-related outcomes (eg, attitudes toward vaccination and intention to get vaccinated [61]; higher presence resulted in more reported cravings for tobacco in [37]). However, presence is rarely measured as a moderator or covariate across studies despite its potential impact on outcomes. The same applies to immersion, which was not measured across studies, despite studies often comparing different intervention modalities of varying degrees of immersion (eg, VR versus 2D screen). VR is not the only technology able to generate presence; narrative, videos, or nonimmersive VR can too [14]. Not measuring immersion or presence across different modalities limits the understanding of VR's role in driving effects on the target outcomes.

Finally, cybersickness was rarely measured across studies despite its potential negative effect on user experience and, in turn, target outcomes. Some studies, notably the ones focusing on physical activity, measured cybersickness and found that it can completely erase the positive effects of using VR (eg, walking on a treadmill while wearing a VR device led to cybersickness, which diminished the positive effects of being exposed to nature compared to the other condition, [50]). Participants who felt symptoms of cybersickness believed that it impacted their experience [59], sometimes to the point they had to drop out of the experiment [62].

## ***Discussion***

### **Principal Findings**

This scoping review identified 51 studies published over the past 14 years that explored the use of VR in health promotion and prevention psychology. Our findings revealed three primary applications of VR: (1) as a tool to deliver interventions, either in feasibility testing or actual implementation; (2) as a means to address fundamental research questions; and (3) as an assessment tool for health-related outcomes. Although VR shows significant promise in creating immersive and engaging interventions, our review highlights the variability in effectiveness and common challenges such as small sample sizes, short follow-up periods, and limited methodological standardization.

VR technology use for health promotion and prevention research is relatively recent, with studies in this review indicating its potential as a promising tool to deliver and assess interventions. For instance, VR was effective in simulating realistic scenarios to engage participants in skills-based learning and decision-making tasks, such as risk-reduction behaviors [42,61]. VR allows researchers to create safe, ecological, and standardized VEs in which it is possible to deliver and evaluate health promotion and preventive interventions [42]; recreate situations or environments that can elicit strong emotional, physiological, behavioral, or psychological responses (eg, mortality salience [29]); and assess outcomes (eg, cravings, food choices) with a multimeasure approach included in VR technologies (eg, psychological, physiological, and behavioral measures). This scoping review identified 51 studies concerning the use of VR technology in the field of health promotion and prevention psychology published within the past 14 years. We mapped (1) the goals and research questions addressed through the use of VR in this field, (2) its effects in the identified areas, and (3) its main challenges or limitations. We identified three main applications of VR in this field: (1) as a tool to deliver an intervention, either (1a) pilot or feasibility testing VR materials or procedures or (1b) using VR to deliver an actual intervention (eg, skills learning, comparing VR vs other intervention modalities) to test its relative efficacy; (2) as a tool to address fundamental research questions; and (3) as an assessment tool.

### **Comparison to Prior Work**

Due to the relative novelty of VR in this field, only 51 eligible studies were published in the past 14 years. Research so far has mostly focused on feasibility or pilot studies, aimed at testing the ability of VR to be integrated into interventions [69], with

a minority of studies focusing on answering fundamental research questions through the use of VR [28]. Most studies employed semiexperimental designs without a control or comparison group and often had a short or no follow-up, limiting the validity and generalizability of results. Studies also included relatively small samples and were often nonrepresentative of the general population (eg, students). However, as the use of VR in the field of health promotion and prevention is still in its infancy, it appears natural to see a stronger focus on pilot or feasibility studies in the published literature.

### Strength and Limitations of the Scoping Review

When considering whether VR is effective in health prevention, it should first be noted that the effectiveness of VR interventions was variable. For instance, while some studies indicated that VR could enhance user engagement and motivation [59], others found no significant differences in outcomes compared to traditional methods [71]. This highlights the need for further research to clarify the conditions under which VR is most effective. This scoping review showed that sometimes VR use is not systematically more effective in achieving target outcomes than its nonimmersive equivalents [41,54,71]; we supposed that to be more effective, VR should be used for its specific immersive characteristics, such as gamification or embodiment, which directly involve the user. For example, skills practice in VR was more effective than role-playing in real life to learn about risk behaviors and ways to avoid them (eg, buying condoms for safer sex) due to VR scenarios' ability to recreate a situation that is realistic, induce emotional changes in the user as the scenario goes on, and finally, make the user have a real first-person experience [42]. Similar results appeared in [61], in which VR was used to represent a vaccination intervention to stop flu spread (ie, participants used handheld controllers to actively send immune cells to prevent flu transmission), whereas in other conditions, participants were just passively watching (a video on a 2D screen or a pamphlet).

It is important to consider the limitations of this review when interpreting the findings. First, as the use of VR in health promotion and prevention psychology is a relatively recent phenomenon, our literature search focused on the last 14 years (2010 - 2024). This resulted in the inclusion of 51 eligible articles, which may have excluded earlier or less accessible studies. However, the majority of included studies (63%) were published between 2020 and 2024, reflecting the increasing affordability and accessibility of VR technology for research in recent years. Therefore, the likelihood of missing pivotal studies is low. Second, the search strategy did not include gray literature, which may have reduced the total number of eligible articles and introduced publication bias by excluding studies with nonsignificant or null results (the file drawer effect). To address this gap, future reviews should consider including gray literature to provide a more comprehensive overview of the field. Third, some studies lacked sufficient reporting of critical aspects such as sample characteristics (eg, size and demographics) and details of VR implementation (eg, exposure duration, type of VR technology used). This limited our ability to draw broad conclusions about the efficacy and applicability of VR in this area. Addressing these reporting gaps in future research will improve the comparability and quality of evidence

in this rapidly evolving area of study. Fourth, although our literature search was updated during the initial revision, which was completed just a few weeks prior to this submission, we recognize that VR research is advancing rapidly. It is therefore possible that new studies may emerge shortly after the conclusion of our search period, which may influence the results of future reviews. To address this, future updates could consider conducting more frequent searches or establishing a continuous review process to ensure that all emerging data are included in real time. However, we are confident that this review accurately reflects the state of the literature as of our latest search.

### Perspectives and Future Research Directions

#### *Standardization of Designs and Replication*

Although the results of our scoping review suggest that VR has potential as a tool for health promotion, the field is still in its infancy. Many studies in this area are limited by small sample sizes, short follow-up periods, and inadequate experimental control. Replication is essential to strengthen the reliability and validity of these findings [72,73]. Replication of these studies in diverse populations and settings will help confirm the generalizability of the findings and identify any boundary conditions, such as differences in user demographics, technology exposure, or the specific health behaviors targeted [74,75]. In addition, replication can shed light on the mechanisms underlying the effectiveness of VR interventions, which may vary depending on the context and population studied. Therefore, further replication is essential not only to solidify current evidence, but also to ensure that VR interventions are applicable and effective across a wide range of health promotion and primary prevention efforts. Replicating existing results to increase the amount and quality of empirical evidence supporting the use and benefit of VR in this field is needed. For example, in this scoping review, we saw that individuals showed an increased knowledge regarding health-related topics [20,69] or changed their behavioral intentions [44,61] when exposed to a VR intervention. However, not all studies provided evidence to fully support these claims [47,63], in addition to the lack of any perspective on how long these effects last or if they are applicable to less specific populations. Therefore, a focus on study replication can strengthen the advancement of research in this field and at the same time prevent a replication crisis, as observed in other fields of behavioral sciences and medicine [76]. There is also a critical need for future research to employ longitudinal study designs. Long-term follow-up is particularly important in preventive psychology, where sustained behavior change and long-term health outcomes are key indicators of success.

#### *VR vs Nonimmersive Apparatus*

The effectiveness of VR compared to nonimmersive interventions, such as 2D presentations, remains controversial. Evidence from the studies included in this review showed mixed results. Although some studies reported that VR interventions increased engagement and enjoyment, others found no significant differences in outcomes compared to nonimmersive methods [59,71]. The immersive features of VR, such as gamification and embodiment, appear to be particularly effective in scenarios that require active user involvement. For example,



participants who practiced risk-avoidance skills in VR showed better retention than those who used real-life role-playing [42]. Similarly, the use of VR in interactive scenarios, such as vaccination education, showed higher levels of engagement than passive modalities such as 2D videos or pamphlets [61]. However, studies have also shown that VR does not always outperform traditional methods in terms of physical activity or knowledge acquisition. This variability highlights the need for future research to clarify the specific contexts in which the immersive qualities of VR are most effective. Systematic assessment of key mechanisms such as presence and immersion could help determine whether VR's effectiveness is primarily due to its immersive nature or to other factors such as interactivity or novelty.

Assessing presence and immersion is crucial for understanding the mechanisms underlying VR and its effects [14]. Evaluating the feeling of presence helps determine the extent to which participants are psychologically immersed in VEs and allows for the identification and correction of potential errors in the VE that could influence presence and, consequently, the effectiveness of VR-based interventions or content. Additionally, addressing such errors can prevent cybersickness and ensure the smooth execution of experiments [50]. Measuring presence and immersion provides valuable insights into individuals' capacity to engage with VR compared to nonimmersive interventions and helps identify how these factors correlate with target outcomes.

### **Set Up for Success**

Conducting feasibility or pilot studies to test the VR procedure and VEs is recommended. As shown in the scoping review, evaluating the enjoyability, usability, and safety of the procedure can be very helpful. Finally, co-designing the VR-based intervention with participants from the targeted population can

enhance the relevance, validity, and user experience with the intervention itself. Cocreating a procedure with participants could induce a bias in their judgment, making them judge the intervention more positively than it actually is. Pilot testing with different groups of participants is recommended to validate the final design.

### **Make It Simple and Clear for Participants**

VR studies are attractive to participants (eg, higher attendance for intervention sessions than the non-VR condition in [19,42,77]), but they can be complex to follow all the way through (ie, risk of cognitive overload, fatigue [78]). When designing studies using VR, keeping them as simple and short as possible will minimize participant burden and fatigue. It is also highly possible that most participants have never experienced VR before, so making sure they understand how to move and interact with the environment at first is necessary. If possible, we recommend doing a short tutorial on how to use the controllers or putting the participant in a tutorial VE before the experimental procedure. The participants can then fully concentrate on what is happening in the VR rather than think about how to interact with the VE.

### **Conclusion**

This scoping review provides an overview of VR's emerging role in health promotion and prevention psychology, highlighting its potential to create immersive and engaging interventions. Although VR has shown promise in delivering health interventions and answering fundamental research questions, its effectiveness remains variable, and many studies are limited by methodological constraints. Future research should prioritize replication, longitudinal designs, and standardized methodologies to strengthen the evidence base and realize the full potential of VR in this field.

---

## **Acknowledgments**

We would like to acknowledge the support of our respective institutions: University Paris Nanterre, Université Clermont Auvergne, Université Rouen Normandie, and Erasmus University Rotterdam. We also thank our colleagues for their insightful discussions and suggestions while conducting this research. This study has been funded by an IresP-INCa grant (number 19II028-00).

---

## **Data Availability**

This scoping review is based on a synthesis of publicly available research articles. The datasets analyzed during this study are derived from published sources that are referenced within the manuscript. No new datasets were generated.

---

## **Authors' Contributions**

SB contributed to the conceptualization, formal analysis, methodology, and writing of the original draft of the manuscript. OZ was involved in the conceptualization, formal analysis, methodology, writing of the original draft, and reviewing and editing the manuscript, while also providing supervision throughout the project. MB participated in the conceptualization, methodology, and writing of the original draft, and offered supervision during the research process.

---

## **Multimedia Appendix 1**

Summary of Studies Using Virtual Reality for Health-Related Behavioral Interventions.

[[DOCX File, 37 KB](#) - [xr\\_v2i1e49923\\_app1.docx](#) ]

---

## **Checklist 1**

PRISMA-ScR checklist.

[[PDF File, 326 KB - xr\\_v2i1e49923\\_app2.pdf](#)]

## References

1. Green LW, Kreuter M. Health Program Planning: An Educational and Ecological Approach, 4th edition: McGraw Hill Higher Education; 2004.
2. McLeroy KR, Bibeau D, Steckler A, Glanz K. An ecological perspective on health promotion programs. *Health Educ Q* 1988;15(4):351-377. [doi: [10.1177/109019818801500401](https://doi.org/10.1177/109019818801500401)] [Medline: [3068205](https://pubmed.ncbi.nlm.nih.gov/3068205/)]
3. Kennedy W, Fruin R, Lue A, Logan SW. Using ecological models of health behavior to promote health care access and physical activity engagement for persons with disabilities. *J Patient Exp* 2021;8:23743735211034031. [doi: [10.1177/23743735211034031](https://doi.org/10.1177/23743735211034031)] [Medline: [34350340](https://pubmed.ncbi.nlm.nih.gov/34350340/)]
4. Schultheis MT, Rizzo AA. The application of virtual reality technology in rehabilitation. *Rehab Psychol* 2001;46(3):296-311. [doi: [10.1037//0090-5550.46.3.296](https://doi.org/10.1037//0090-5550.46.3.296)]
5. Ahn SJG. Incorporating immersive virtual environments in health promotion campaigns: a construal level theory approach. *Health Commun* 2015;30(6):545-556. [doi: [10.1080/10410236.2013.869650](https://doi.org/10.1080/10410236.2013.869650)] [Medline: [24991725](https://pubmed.ncbi.nlm.nih.gov/24991725/)]
6. Bailey JO, Bailenson JN, Flora J, Armel KC, Voelker D, Reeves B. The impact of vivid messages on reducing energy consumption related to hot water use. *Env Behav* 2015 Jun;47(5):570-592. [doi: [10.1177/0013916514551604](https://doi.org/10.1177/0013916514551604)]
7. Bailenson J. Experience on Demand: What Virtual Reality Is, How It Works, and What It Can Do: WW Norton & Company; 2018. URL: <https://psycnet.apa.org/record/2019-04570-000> [accessed 2024-12-15]
8. Rizzo A, Bouchard S. Virtual Reality for Psychological and Neurocognitive Interventions: Springer; 2019.
9. McCall C, Blascovich J. How, when, and why to use digital experimental virtual environments to study social behavior. *Soc & Pers Psych* 2009 Sep;3(5):744-758. [doi: [10.1111/j.1751-9004.2009.00195.x](https://doi.org/10.1111/j.1751-9004.2009.00195.x)]
10. Yaremchuk HE, Persky S. Tracing physical behavior in virtual reality: a narrative review of applications to social psychology. *J Exp Soc Psychol* 2019 Nov;85:103845. [doi: [10.1016/j.jesp.2019.103845](https://doi.org/10.1016/j.jesp.2019.103845)] [Medline: [32831397](https://pubmed.ncbi.nlm.nih.gov/32831397/)]
11. Cummings JJ, Bailenson JN. How immersive is enough? A meta-analysis of the effect of immersive technology on user presence. *Med Psychol* 2016 Apr 2;19(2):272-309. [doi: [10.1080/15213269.2015.1015740](https://doi.org/10.1080/15213269.2015.1015740)]
12. Parsons TD, Gaggioli A, Riva G. Virtual reality for research in social neuroscience. *Brain Sci* 2017 Apr 16;7(4):42. [doi: [10.3390/brainsci7040042](https://doi.org/10.3390/brainsci7040042)] [Medline: [28420150](https://pubmed.ncbi.nlm.nih.gov/28420150/)]
13. Ahn SJG, Hahm JM, Johnsen K. Feeling the weight of calories: using haptic feedback as virtual exemplars to promote risk perception among young females on unhealthy snack choices. *Med Psychol* 2019;22(4):626-652. [doi: [10.1080/15213269.2018.1492939](https://doi.org/10.1080/15213269.2018.1492939)] [Medline: [32863775](https://pubmed.ncbi.nlm.nih.gov/32863775/)]
14. Pianzola F, Riva G, Mantovani F, Kukkonen K. Presence, flow, and narrative absorption: an interdisciplinary theoretical exploration with a new spatiotemporal integrated model based on predictive processing. *PsyArXiv*. Preprint posted online on Jul 2, 2020. [doi: [10.31234/osf.io/uzrcm](https://doi.org/10.31234/osf.io/uzrcm)]
15. Yao S, Kim G. The effects of immersion in a virtual reality game: presence and physical activity. Presented at: 21st International Conference on Human-Computer Interaction; Jul 26-31, 2019; Orlando, FL, USA p. 234-242. [doi: [10.1007/978-3-030-22602-2\\_18](https://doi.org/10.1007/978-3-030-22602-2_18)]
16. Schwind V, Knierim P, Haas N, Henze N. Using presence questionnaires in virtual reality. Presented at: Proceedings of the 2019 CHI Conference on Human Factors in Computing Systems; May 2, 2019; Glasgow, Scotland, UK p. 1-12. [doi: [10.1145/3290605.3300590](https://doi.org/10.1145/3290605.3300590)]
17. Bonnetterre S, Zerhouni O, Boffo M. The influence of billboard-based tobacco prevention posters on memorization, attitudes, and craving: immersive virtual reality study. *J Med Internet Res* 2024 Jul 9;26:e49344. [doi: [10.2196/49344](https://doi.org/10.2196/49344)] [Medline: [38980707](https://pubmed.ncbi.nlm.nih.gov/38980707/)]
18. Schmäzle R, Lim S, Cho HJ, Wu J, Bente G. Examining the exposure-reception-retention link in realistic communication environments via VR and eye-tracking: the VR billboard paradigm. *PLoS One* 2023;18(11):e0291924. [doi: [10.1371/journal.pone.0291924](https://doi.org/10.1371/journal.pone.0291924)] [Medline: [38033032](https://pubmed.ncbi.nlm.nih.gov/38033032/)]
19. Riva G, Baños RM, Botella C, Mantovani F, Gaggioli A. Transforming experience: the potential of augmented reality and virtual reality for enhancing personal and clinical change. *Front Psychiatry* 2016;7:164. [doi: [10.3389/fpsy.2016.00164](https://doi.org/10.3389/fpsy.2016.00164)] [Medline: [27746747](https://pubmed.ncbi.nlm.nih.gov/27746747/)]
20. Saab MM, Landers M, Cooke E, Murphy D, Davoren M, Hegarty J. Enhancing men's awareness of testicular disorders using a virtual reality intervention. *Nurs Res* 2018;67(5):349-358. [doi: [10.1097/NNR.0000000000000303](https://doi.org/10.1097/NNR.0000000000000303)] [Medline: [30059354](https://pubmed.ncbi.nlm.nih.gov/30059354/)]
21. Hamilton-Giachritsis C, Banakou D, Garcia Quiroga M, Giachritsis C, Slater M. Reducing risk and improving maternal perspective-taking and empathy using virtual embodiment. *Sci Rep* 2018 Feb 14;8(1):2975. [doi: [10.1038/s41598-018-21036-2](https://doi.org/10.1038/s41598-018-21036-2)] [Medline: [29445183](https://pubmed.ncbi.nlm.nih.gov/29445183/)]
22. Tricco AC, Lillie E, Zarin W, et al. A scoping review on the conduct and reporting of scoping reviews. *BMC Med Res Methodol* 2016 Feb 9;16:15. [doi: [10.1186/s12874-016-0116-4](https://doi.org/10.1186/s12874-016-0116-4)] [Medline: [26857112](https://pubmed.ncbi.nlm.nih.gov/26857112/)]
23. Tricco AC, Lillie E, Zarin W, et al. PRISMA extension for Scoping Reviews (PRISMA-ScR): checklist and explanation. *Ann Intern Med* 2018 Oct 2;169(7):467-473. [doi: [10.7326/M18-0850](https://doi.org/10.7326/M18-0850)] [Medline: [30178033](https://pubmed.ncbi.nlm.nih.gov/30178033/)]



24. Campbell M, McKenzie JE, Sowden A, et al. Synthesis without meta-analysis (SWiM) in systematic reviews: reporting guideline. *BMJ* 2020 Jan 16;368:16890. [doi: [10.1136/bmj.16890](https://doi.org/10.1136/bmj.16890)] [Medline: [31948937](https://pubmed.ncbi.nlm.nih.gov/31948937/)]
25. Lemieux V, Monthuy-Blanc J, Corno G. Application of virtual reality in an ED primary prevention context: an exploratory study. *Annu Rev CyberTher Telemed* 2020;18:207-211 [FREE Full text]
26. Blom S, Gillebaart M, De Boer F, van der Laan N, De Ridder DTD. Under pressure: nudging increases healthy food choice in a virtual reality supermarket, irrespective of system 1 reasoning. *Appetite* 2021 May 1;160:105116. [doi: [10.1016/j.appet.2021.105116](https://doi.org/10.1016/j.appet.2021.105116)] [Medline: [33450297](https://pubmed.ncbi.nlm.nih.gov/33450297/)]
27. Isgin-Atici K, Ozkan A, Celikcan U, et al. Usability study of a novel tool: the virtual cafeteria in nutrition education. *J Nutr Educ Behav* 2020 Nov;52(11):1058-1065. [doi: [10.1016/j.jneb.2020.08.001](https://doi.org/10.1016/j.jneb.2020.08.001)] [Medline: [33168205](https://pubmed.ncbi.nlm.nih.gov/33168205/)]
28. Ledoux T, Nguyen AS, Bakos-Block C, Bordnick P. Using virtual reality to study food cravings. *Appetite* 2013 Dec;71:396-402. [doi: [10.1016/j.appet.2013.09.006](https://doi.org/10.1016/j.appet.2013.09.006)] [Medline: [24055758](https://pubmed.ncbi.nlm.nih.gov/24055758/)]
29. Marcum CS, Goldring MR, McBride CM, Persky S. Modeling dynamic food choice processes to understand dietary intervention effects. *Ann Behav Med* 2018 Feb 17;52(3):252-261. [doi: [10.1093/abm/kax041](https://doi.org/10.1093/abm/kax041)] [Medline: [29538666](https://pubmed.ncbi.nlm.nih.gov/29538666/)]
30. McBride CM, Persky S, Wagner LK, Faith MS, Ward DS. Effects of providing personalized feedback of child's obesity risk on mothers' food choices using a virtual reality buffet. *Int J Obes (Lond)* 2013 Oct;37(10):1322-1327. [doi: [10.1038/ijo.2013.87](https://doi.org/10.1038/ijo.2013.87)] [Medline: [23736369](https://pubmed.ncbi.nlm.nih.gov/23736369/)]
31. Persky S, Ferrer RA, Klein WMP, et al. Effects of fruit and vegetable feeding messages on mothers and fathers: interactions between emotional state and health message framing. *Ann Behav Med* 2019 Aug 16;53(9):789-800. [doi: [10.1093/abm/kay088](https://doi.org/10.1093/abm/kay088)] [Medline: [30395145](https://pubmed.ncbi.nlm.nih.gov/30395145/)]
32. Persky S, Yaremych HE, Goldring MR, Ferrer RA, Rose MK, Hollister BM. Investigating the efficacy of genetic, environmental, and multifactorial risk information when communicating obesity risk to parents of young children. *Ann Behav Med* 2021 Jul 22;55(8):720-733. [doi: [10.1093/abm/kaaa103](https://doi.org/10.1093/abm/kaaa103)] [Medline: [33196082](https://pubmed.ncbi.nlm.nih.gov/33196082/)]
33. Verhulst A, Normand JM, Lombart C, Sugimoto M, Moreau G. Influence of being embodied in an obese virtual body on shopping behavior and products perception in VR. *Front Robot AI* 2018;5:113. [doi: [10.3389/frobt.2018.00113](https://doi.org/10.3389/frobt.2018.00113)] [Medline: [33500992](https://pubmed.ncbi.nlm.nih.gov/33500992/)]
34. Ferrer-Garcia M, Gutiérrez-Maldonado J, Agliaro-López M, Lobera-Espi X, Pla J, Vilalta-Abella F. Validation of VR-based software for binge eating treatment: preliminary data. *Annu Rev CyberTher Telemed* 2014;12:146-150. [doi: [10.3233/978-1-61499-401-5-146](https://doi.org/10.3233/978-1-61499-401-5-146)]
35. Borrelli B, Ruelas N, Jurassic M. Delivery of a smoking cessation induction intervention via virtual reality headset during a dental cleaning. *Transl Behav Med* 2021 Feb 11;11(1):182-188. [doi: [10.1093/tbm/ibz144](https://doi.org/10.1093/tbm/ibz144)] [Medline: [31665495](https://pubmed.ncbi.nlm.nih.gov/31665495/)]
36. Ferrer-García M, García-Rodríguez O, Gutiérrez-Maldonado J, Pericot-Valverde I, Secades-Villa R. Efficacy of virtual reality in triggering the craving to smoke: its relation to level of presence and nicotine dependence. *Annu Rev CyberTher Telemed* 2010;8:99-102. [doi: [10.3233/978-1-60750-561-7-123](https://doi.org/10.3233/978-1-60750-561-7-123)]
37. García-Rodríguez O, Weidberg S, Gutiérrez-Maldonado J, Secades-Villa R. Smoking a virtual cigarette increases craving among smokers. *Addict Behav* 2013 Oct;38(10):2551-2554. [doi: [10.1016/j.addbeh.2013.05.007](https://doi.org/10.1016/j.addbeh.2013.05.007)] [Medline: [23793042](https://pubmed.ncbi.nlm.nih.gov/23793042/)]
38. Weser VU, Duncan LR, Pendergrass TM, Fernandes CS, Fiellin LE, Hieftje KD. A quasi-experimental test of a virtual reality game prototype for adolescent e-cigarette prevention. *Addict Behav* 2021 Jan;112:106639. [doi: [10.1016/j.addbeh.2020.106639](https://doi.org/10.1016/j.addbeh.2020.106639)] [Medline: [32920455](https://pubmed.ncbi.nlm.nih.gov/32920455/)]
39. Weser VU, Duncan LR, Sands BE, et al. Evaluation of a virtual reality e-cigarette prevention game for adolescents. *Addict Behav* 2021 Nov;122:107027. [doi: [10.1016/j.addbeh.2021.107027](https://doi.org/10.1016/j.addbeh.2021.107027)] [Medline: [34225030](https://pubmed.ncbi.nlm.nih.gov/34225030/)]
40. Guldager JD, Kjær SL, Grittner U, Stock C. Efficacy of the virtual reality intervention VR FestLab on alcohol refusal self-efficacy: a cluster-randomized controlled trial. *Int J Environ Res Public Health* 2022 Mar 10;19(6):3293. [doi: [10.3390/ijerph19063293](https://doi.org/10.3390/ijerph19063293)] [Medline: [35328977](https://pubmed.ncbi.nlm.nih.gov/35328977/)]
41. Ma Z. The use of immersive stories to influence college students' attitudes and intentions related to drinking and driving. *J Am Coll Health* 2022 Oct;70(7):2042-2049. [doi: [10.1080/07448481.2020.1842418](https://doi.org/10.1080/07448481.2020.1842418)] [Medline: [33151847](https://pubmed.ncbi.nlm.nih.gov/33151847/)]
42. Hadley W, Houck CD, Barker DH, et al. Eliciting affect via immersive virtual reality: a tool for adolescent risk reduction. *J Pediatr Psychol* 2014 Apr;39(3):358-368. [doi: [10.1093/jpepsy/jst092](https://doi.org/10.1093/jpepsy/jst092)] [Medline: [24365699](https://pubmed.ncbi.nlm.nih.gov/24365699/)]
43. Hadley W, Houck C, Brown LK, Spitalnick JS, Ferrer M, Barker D. Moving beyond role-play: evaluating the use of virtual reality to teach emotion regulation for the prevention of adolescent risk behavior within a randomized pilot trial. *J Pediatr Psychol* 2019 May 1;44(4):425-435. [doi: [10.1093/jpepsy/jsy092](https://doi.org/10.1093/jpepsy/jsy092)] [Medline: [30551157](https://pubmed.ncbi.nlm.nih.gov/30551157/)]
44. Detez L, Greenwood LM, Segrave R, et al. A psychophysiological and behavioural study of slot machine near-misses using immersive virtual reality. *J Gambl Stud* 2019 Sep;35(3):929-944. [doi: [10.1007/s10899-018-09822-z](https://doi.org/10.1007/s10899-018-09822-z)] [Medline: [30684139](https://pubmed.ncbi.nlm.nih.gov/30684139/)]
45. Alyan E, Combe T, Awang Rambli DR, Sulaiman S, Merienne F, Muhaiyuddin NDM. The influence of virtual forest walk on physiological and psychological responses. *Int J Environ Res Public Health* 2021 Oct 29;18(21):11420. [doi: [10.3390/ijerph182111420](https://doi.org/10.3390/ijerph182111420)] [Medline: [34769937](https://pubmed.ncbi.nlm.nih.gov/34769937/)]
46. Beverly E, Hommema L, Coates K, et al. A tranquil virtual reality experience to reduce subjective stress among COVID-19 frontline healthcare workers. *PLoS ONE* 2022;17(2):e0262703. [doi: [10.1371/journal.pone.0262703](https://doi.org/10.1371/journal.pone.0262703)] [Medline: [35139092](https://pubmed.ncbi.nlm.nih.gov/35139092/)]

47. Brimelow RE, Dawe B, Dissanayaka N. Preliminary research: virtual reality in residential aged care to reduce apathy and improve mood. *Cyberpsychol Behav Soc Netw* 2020 Mar;23(3):165-170. [doi: [10.1089/cyber.2019.0286](https://doi.org/10.1089/cyber.2019.0286)] [Medline: [31829729](https://pubmed.ncbi.nlm.nih.gov/31829729/)]
48. Brimelow RE, Thangavelu K, Beattie R, Dissanayaka NN. Feasibility of group-based multiple virtual reality sessions to reduce behavioral and psychological symptoms in persons living in residential aged care. *J Am Med Dir Assoc* 2022 May;23(5):831-837. [doi: [10.1016/j.jamda.2021.07.026](https://doi.org/10.1016/j.jamda.2021.07.026)] [Medline: [34454923](https://pubmed.ncbi.nlm.nih.gov/34454923/)]
49. Browning M, Mimnaugh KJ, van Riper CJ, Laurent HK, LaValle SM. Can simulated nature support mental health? Comparing short, single-doses of 360-degree nature videos in virtual reality with the outdoors. *Front Psychol* 2019;10:2667. [doi: [10.3389/fpsyg.2019.02667](https://doi.org/10.3389/fpsyg.2019.02667)] [Medline: [32010003](https://pubmed.ncbi.nlm.nih.gov/32010003/)]
50. Calogiuri G, Litleskare S, Fagerheim KA, Rydgren TL, Brambilla E, Thurston M. Experiencing nature through immersive virtual environments: environmental perceptions, physical engagement, and affective responses during a simulated nature walk. *Front Psychol* 2017;8:2321. [doi: [10.3389/fpsyg.2017.02321](https://doi.org/10.3389/fpsyg.2017.02321)] [Medline: [29410635](https://pubmed.ncbi.nlm.nih.gov/29410635/)]
51. Afifi T, Collins N, Rand K, et al. Using virtual reality to improve the quality of life of older adults with cognitive impairments and their family members who live at a distance. *Health Commun* 2023 Oct;38(9):1904-1915. [doi: [10.1080/10410236.2022.2040170](https://doi.org/10.1080/10410236.2022.2040170)] [Medline: [35253531](https://pubmed.ncbi.nlm.nih.gov/35253531/)]
52. Adhyaru JS, Kemp C. Virtual reality as a tool to promote wellbeing in the workplace. *D Health* 2022;8:20552076221084473. [doi: [10.1177/20552076221084473](https://doi.org/10.1177/20552076221084473)] [Medline: [35284084](https://pubmed.ncbi.nlm.nih.gov/35284084/)]
53. Kim J, Jung YH, Shin YB, et al. Development and validation of a virtual reality-based training program for promoting subjective well-being. *Psychiatry Investig* 2020 Dec;17(12):1207-1215. [doi: [10.30773/pi.2020.0311](https://doi.org/10.30773/pi.2020.0311)] [Medline: [33301665](https://pubmed.ncbi.nlm.nih.gov/33301665/)]
54. Riva G, Bernardelli L, Castelnuovo G, et al. A virtual reality-based self-help intervention for dealing with the psychological distress associated with the COVID-19 lockdown: an effectiveness study with a two-week follow-up. *Int J Environ Res Public Health* 2021 Aug 2;18(15):8188. [doi: [10.3390/ijerph18158188](https://doi.org/10.3390/ijerph18158188)] [Medline: [34360479](https://pubmed.ncbi.nlm.nih.gov/34360479/)]
55. Ko H, Lim HJ, Park J, et al. Subjective experience on virtual reality-assisted mental health promotion program. *Psychiatry Investig* 2024 Apr;21(4):380-386. [doi: [10.30773/pi.2023.0286](https://doi.org/10.30773/pi.2023.0286)] [Medline: [38695045](https://pubmed.ncbi.nlm.nih.gov/38695045/)]
56. Kiper P, Przysiężna E, Cieślak B, et al. Effects of immersive virtual therapy as a method supporting recovery of depressive symptoms in post-stroke rehabilitation: randomized controlled trial. *Clin Interv Aging* 2022;17:1673-1685. [doi: [10.2147/CIA.S375754](https://doi.org/10.2147/CIA.S375754)] [Medline: [36447623](https://pubmed.ncbi.nlm.nih.gov/36447623/)]
57. Eisapour M, Cao S, Boger J. Participatory design and evaluation of virtual reality games to promote engagement in physical activity for people living with dementia. *J Rehabil Assist Technol Eng* 2020;7:2055668320913770. [doi: [10.1177/2055668320913770](https://doi.org/10.1177/2055668320913770)] [Medline: [32499921](https://pubmed.ncbi.nlm.nih.gov/32499921/)]
58. Fang YM, Huang YJ. Comparison of the usability and flow experience of an exercise promotion virtual reality programme for different age groups. *Behav Inf Technol* 2021 Sep 10;40(12):1250-1264. [doi: [10.1080/0144929X.2021.1938680](https://doi.org/10.1080/0144929X.2021.1938680)]
59. Farič N, Smith L, Hon A, et al. A virtual reality exergame to engage adolescents in physical activity: mixed methods study describing the formative intervention development process. *J Med Internet Res* 2021 Feb 4;23(2):e18161. [doi: [10.2196/18161](https://doi.org/10.2196/18161)] [Medline: [33538697](https://pubmed.ncbi.nlm.nih.gov/33538697/)]
60. Bird JM, Karageorghis CI, Baker SJ, Brookes DA, Nowicky AV. Ready Exerciser One: effects of music and virtual reality on cycle ergometer exercise. *Br J Health Psychol* 2021 Feb;26(1):15-32. [doi: [10.1111/bjhp.12445](https://doi.org/10.1111/bjhp.12445)] [Medline: [32538512](https://pubmed.ncbi.nlm.nih.gov/32538512/)]
61. Zeng N, Liu W, Pope ZC, McDonough DJ, Gao Z. Acute effects of virtual reality exercise biking on college students' physical responses. *Res Q Exerc Sport* 2022 Sep;93(3):633-639. [doi: [10.1080/02701367.2021.1891188](https://doi.org/10.1080/02701367.2021.1891188)] [Medline: [34663191](https://pubmed.ncbi.nlm.nih.gov/34663191/)]
62. Zeng N, Pope Z, Gao Z. Acute effect of virtual reality exercise bike games on college students' physiological and psychological outcomes. *Cyberpsychol Behav Soc Netw* 2017 Jul;20(7):453-457. [doi: [10.1089/cyber.2017.0042](https://doi.org/10.1089/cyber.2017.0042)] [Medline: [28715263](https://pubmed.ncbi.nlm.nih.gov/28715263/)]
63. Mottelson A, Vandeweerd C, Atchapero M, et al. A self-administered virtual reality intervention increases COVID-19 vaccination intention. *Vaccine* 2021 Nov 5;39(46):6746-6753. [doi: [10.1016/j.vaccine.2021.10.004](https://doi.org/10.1016/j.vaccine.2021.10.004)] [Medline: [34654579](https://pubmed.ncbi.nlm.nih.gov/34654579/)]
64. Nowak GJ, Evans NJ, Wojdyski BW, et al. Using immersive virtual reality to improve the beliefs and intentions of influenza vaccine avoidant 18-to-49-year-olds: considerations, effects, and lessons learned. *Vaccine* 2020 Jan 29;38(5):1225-1233. [doi: [10.1016/j.vaccine.2019.11.009](https://doi.org/10.1016/j.vaccine.2019.11.009)] [Medline: [31806533](https://pubmed.ncbi.nlm.nih.gov/31806533/)]
65. Niki K, Yasui M, Iguchi M, Isono T, Kageyama H, Ueda M. A pilot study to develop a new method of assisting children in taking their medication by using immersive virtual reality. *Biol Pharm Bull* 2021;44(2):279-282. [doi: [10.1248/bpb.b20-00646](https://doi.org/10.1248/bpb.b20-00646)] [Medline: [33518682](https://pubmed.ncbi.nlm.nih.gov/33518682/)]
66. Polechoński J, Nierwińska K, Kalita B, Wodarski P. Can physical activity in immersive virtual reality be attractive and have sufficient intensity to meet health recommendations for obese children? A pilot study. *Int J Environ Res Public Health* 2020 Nov 1;17(21):8051. [doi: [10.3390/ijerph17218051](https://doi.org/10.3390/ijerph17218051)] [Medline: [33139604](https://pubmed.ncbi.nlm.nih.gov/33139604/)]
67. Sakhare A, Stradford J, Ravichandran R, et al. Simultaneous exercise and cognitive training in virtual reality phase 2 pilot study: impact on brain health and cognition in older adults. *Brain Plast* 2021;7(2):111-130. [doi: [10.3233/BPL-210126](https://doi.org/10.3233/BPL-210126)] [Medline: [34868877](https://pubmed.ncbi.nlm.nih.gov/34868877/)]
68. Horsham C, Dutton-Regester K, Antrobus J, et al. A virtual reality game to change sun protection behavior and prevent cancer: user-centered design approach. *JMIR Serious Games* 2021 Mar 25;9(1):e24652. [doi: [10.2196/24652](https://doi.org/10.2196/24652)] [Medline: [33764308](https://pubmed.ncbi.nlm.nih.gov/33764308/)]

69. Guo JL, Hsu HP, Lai TM, Lin ML, Chung CM, Huang CM. Acceptability evaluation of the use of virtual reality games in smoking-prevention education for high school students: prospective observational study. *J Med Internet Res* 2021 Sep 28;23(9):e28037. [doi: [10.2196/28037](https://doi.org/10.2196/28037)] [Medline: [34581679](https://pubmed.ncbi.nlm.nih.gov/34581679/)]
70. Chittaro L, Sioni R, Crescentini C, Fabbro F. Mortality salience in virtual reality experiences and its effects on users' attitudes towards risk. *Int J Hum Comput Stud* 2017 May;101:10-22. [doi: [10.1016/j.ijhcs.2017.01.002](https://doi.org/10.1016/j.ijhcs.2017.01.002)]
71. Namkoong K, Chen J, Leach J, et al. Virtual reality for public health: a study on a VR intervention to enhance occupational injury prevention. *J Public Health (Bangkok)* 2023 Mar 14;45(1):136-144. [doi: [10.1093/pubmed/fdab407](https://doi.org/10.1093/pubmed/fdab407)]
72. Pashler H, Harris CR. Is the replicability crisis overblown? Three arguments examined. *Perspect Psychol Sci* 2012 Nov;7(6):531-536. [doi: [10.1177/1745691612463401](https://doi.org/10.1177/1745691612463401)] [Medline: [26168109](https://pubmed.ncbi.nlm.nih.gov/26168109/)]
73. Schmidt S. Shall we really do it again? The powerful concept of replication is neglected in the social sciences. *Rev Gen Psychol* 2009 Jun;13(2):90-100. [doi: [10.1037/a0015108](https://doi.org/10.1037/a0015108)]
74. Open Science Collaboration. Estimating the reproducibility of psychological science. *Science* 2015 Aug 28;349(6251). [doi: [10.1126/science.aac4716](https://doi.org/10.1126/science.aac4716)]
75. Klein SB. What can recent replication failures tell us about the theoretical commitments of psychology? *Theory Psychol* 2014 Jun;24(3):326-338. [doi: [10.1177/0959354314529616](https://doi.org/10.1177/0959354314529616)]
76. Ahn S. Virtual exemplars in health promotion campaigns: heightening perceived risk and involvement to reduce soft drink consumption in young adults. *J Media Psychol* 2018 Apr;30(2):91-103. [doi: [10.1027/1864-1105/a000184](https://doi.org/10.1027/1864-1105/a000184)]
77. Ahn SJ, Fox J, Dale KR, Avant JA. Framing virtual experiences: effects on environmental efficacy and behavior over time. *Comm Res* 2015;42(6):839-863. [doi: [10.1177/0093650214534973](https://doi.org/10.1177/0093650214534973)]
78. Behr KM, Nosper A, Klimmt C, Hartmann T. Some practical considerations of ethical issues in VR research. *Pres Teleoperators & Virtual Environ* 2005 Dec;14(6):668-676. [doi: [10.1162/105474605775196535](https://doi.org/10.1162/105474605775196535)]

## Abbreviations

**HMD:** head-mounted display

**PRISMA-ScR:** Preferred Reporting Items for Systematic Reviews and Meta-analyses extension for Scoping Reviews

**VE:** virtual environment

**VR:** virtual reality

*Edited by L Riedemann, T Leung; submitted 13.06.23; peer-reviewed by B Jacob, B Cieřlik, L Mendes; revised version received 27.11.24; accepted 27.11.24; published 20.01.25.*

*Please cite as:*

*Bonnetterre S, Zerhouni O, Boffo M*

*Immersive Virtual Reality for Health Promotion and Primary Prevention in Psychology: Scoping Review*

*JMIR XR Spatial Comput* 2025;2:e49923

*URL:* <https://xr.jmir.org/2025/1/e49923>

*doi:* [10.2196/49923](https://doi.org/10.2196/49923)

© Solenne Bonnetterre, Oulmann Zerhouni, Marilisa Boffo. Originally published in JMIR XR and Spatial Computing (<https://xr.jmir.org>), 20.1.2025. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in JMIR XR and Spatial Computing, is properly cited. The complete bibliographic information, a link to the original publication on <https://xr.jmir.org/>, as well as this copyright and license information must be included.

# Augmented Reality in Enhancing Operating Room Crisis Checklist Adherence: Randomized Comparative Efficacy Study

Rayan Ebnali Harari<sup>1\*</sup>, PhD; Abdullah Altaweel<sup>1,2\*</sup>, MD; Erik Anderson<sup>3\*</sup>, MD; Charles Pozner<sup>4\*</sup>, MD; Rafael Grossmann<sup>5\*</sup>, MD; Andrew Goldsmith<sup>3</sup>, MBA, MD; Hamid Shokoohi<sup>3\*</sup>, MD

<sup>1</sup>STRATUS Center for Medical Simulation, Mass General Brigham, Harvard Medical School, 10 Vining, Boston, MA, United States

<sup>2</sup>Kuwait Ministry of Health, Kuwait City, Kuwait

<sup>3</sup>Department of Emergency Medicine, Mass General Brigham, Boston, MA, United States

<sup>4</sup>PC Institute for Medical Education, Boston, MA, United States

<sup>5</sup>Department of Surgery, Portsmouth Regional Hospital, Boston, MA, United States

\* these authors contributed equally

## Corresponding Author:

Rayan Ebnali Harari, PhD

STRATUS Center for Medical Simulation, Mass General Brigham, Harvard Medical School, 10 Vining, Boston, MA, United States

## Abstract

**Background:** Effective crisis management in operating rooms (ORs) is crucial for patient safety. Despite their benefits, adherence to OR crisis checklists is often limited, highlighting the need for innovative solutions.

**Objective:** The objective of this study was to evaluate the efficacy of augmented reality (AR)-enhanced checklists in improving protocol adherence, compared to traditional paper checklists and no checklist scenarios during simulated OR crises.

**Methods:** This study was a randomized comparative efficacy study comparing the utility of AR checklists, paper checklists, and no checklist scenarios using 4 validated and simulated OR crises scenarios: asystolic cardiac arrest, air embolism, unexplained hypotension/hypoxia, and malignant hyperthermia. The study took place in a simulated OR setting and had applicability to the standard procedures in ORs, critical care units, and urgent care scenarios in the emergency department. To form the 24 OR teams, 50 professionals including 24 anesthesiologists, 24 nurses, 1 surgeon, and 1 scrub nurse from two academic hospitals were included. The primary outcome measured was the failure to adhere (FTA) rate for critical actions during simulated OR crises. Adherence was determined using retrospective video analysis involving 595 key processes evaluated across 24 surgical teams. Interrater reliability was assessed using a Cohen  $\kappa$ . Secondary outcomes included checklist usability and cognitive load, as measured by the low-frequency to high-frequency (LF/HF) ratio of the heart rate variability.

**Results:** The AR checklist group showed a significantly lower FTA rate (mean 15.1%, SD 5.77%) compared to the paper checklist (mean 8.32%, SD 5.65%;  $t_{23}=-2.08$ ;  $P=.048$ ) and the no checklist groups (mean 29.81%, SD 5.59%;  $t_{23}=-6.47$ ;  $P<.001$ ). The AR checklist also resulted in a higher LF/HF ratio for anesthesiologists ( $F_{2,46}=4.88$ ;  $P=.02$ ), showing a potential increase in the level of cognitive load. Survey data indicated positive receptions for both AR and paper checklists.

**Conclusions:** These results suggest that AR checklists could offer a viable method for enhancing adherence to critical care protocols. Although, further research is needed to fully assess their impact on clinical outcomes and to address any associated increase in cognitive load.

(JMIR XR Spatial Comput 2025;2:e60792) doi:[10.2196/60792](https://doi.org/10.2196/60792)

## KEYWORDS

augmented reality; operating room; crisis checklist; checklist; guideline adherence; quality improvement; patient safety; cardiac arrest; hypotension; hyperthermia; critical care; emergency department

## Introduction

Unexpected crises in the operating room (OR), such as cardiac arrests or severe hemorrhages, create a critical situation in which surgical teams should deliver rapid and coordinated care with a time-sensitive order of actions listed in the OR crisis checklists

[1-3]. Although these high-stakes, low-frequency crises may occur infrequently for any single practitioner, their cumulative incidence across hospitals underscores a significant challenge to patient safety and surgical outcomes [4-7]. The OR teams' ability to effectively manage these life-threatening complications depends on their preparedness in managing crises [8,9], training [10], and adherence to the validated crisis checklists [11].



Presurgical checklists are used before surgery to ensure correct patient identification and procedure planning. In contrast, crisis management checklists guide surgical teams during emergencies, helping them respond quickly to life-threatening situations. While both checklists improve safety, this study focuses specifically on crisis management checklists, which aim to support decision-making during critical events in the OR.

The lack of adherence to the checklists negatively impacts surgical mortality rates and overall hospital performance [12]. Evidence suggests that adherence to established best practices during these critical moments is varied and often associated with a decay in the retention of essential skills and knowledge over time [13-16]. In many instances, the use of surgical safety checklists was associated with a reduction in morbidity and mortality, and they were integrated as a new standard of care [17,18]. The dynamic and high-pressure nature of surgical emergencies requires not only adherence to protocols but also the ability to quickly access and use complex information under cognitively demanding conditions [19-21]. However, even though adherence to these checklists is crucial, the traditional paper ones are often difficult to use effectively in such intense scenarios [22-24]. The low adoption of checklists underscores the need for innovative approaches to using checklists that fit with surgical workflows, enhancing protocol adherence without disrupting the clinical focus.

Augmented reality (AR) technology, by relaying important procedural information directly into the clinicians' vision [25-28], can enhance protocol adherence in medical settings [29-33]. Initial applications of AR in medication management and emergency trauma care have shown promise in reducing errors and guiding clinicians through complex procedures with enhanced clarity and efficiency [34-38]. This evidence positions AR as a potential technology for improving adherence to

medical protocols [39-41]. However, the effectiveness of and adherence to AR-enhanced surgical checklists during OR crises has not been thoroughly studied.

This study aims to evaluate the efficacy of AR-enhanced checklists in improving protocol adherence by surgical teams during simulated OR crises. By comparing outcomes with the traditional paper checklists and scenarios without a checklist, the research seeks to provide evidence on AR's utility to reduce the failure to adhere (FTA) rate for crucial procedural steps when managing surgical crises, ultimately improving patient outcomes in the OR. We hypothesize that the AR-enhanced checklists will significantly reduce the FTA rate for crucial procedural steps compared to traditional paper checklists and no checklist scenarios.

## Methods

### Study Design


This prospective within-subject study aimed to compare the impact of AR checklists, traditional paper checklists, and no checklist conditions on managing OR crises (Figure 1). A detailed outline of team participation and the methodological framework is included in Multimedia Appendix 1. The development and rationale behind the crisis checklists, guided by surgical safety standards, have been detailed in a previous publication [14]. Teams, including anesthesia staff, OR nurses, and a mock surgeon, faced simulated intraoperative crises with randomized scenario assignments and checklist types. Before the main investigation, a pilot study tested the scenario fidelity and the AR checklist's practicality. Paper checklists were provided in booklet form and placed near the anesthesia machine and the circulating nurse's station, mirroring their accessibility in actual ORs. A summary and the checklists are available in sections 1 - 3 of Multimedia Appendix 1.

**Figure 1.** Study overview diagram. (a) Checklists presented in an augmented reality interface using Microsoft HoloLens 2. (b) Study design scenarios including an augmented reality checklist, paper checklist, and no checklist.

**a**



**b**



**Setups: The OR Checklists**

We used OR crisis checklists for 4 critical scenarios: (1) asystolic cardiac arrest, (2) air embolism, (3) unexplained hypotension/hypoxia, and (4) malignant hyperthermia. These scenarios were derived from a comprehensive checklist development and testing process explained by Ziewacz et al [42] and were chosen for their clinical importance and feasibility for implementation in AR. Additionally, we followed the standardized approach used by Arriaga et al [14], which evaluated the efficacy of these checklists in improving adherence to lifesaving protocols through high-fidelity medical simulations. More details on the checklists and key processes evaluated to measure adherence to protocols can be found in section 3 of [Multimedia Appendix 1](#).

**Participants**

Participants were recruited from 2 academic hospitals between October 2021, and September 2023. Each team comprised the anesthesia staff (including attending physicians and residents), OR nurses, one mock surgeon, and one scrub nurse, totaling 24 attending physicians and residents, 24 OR nurses, and one mock surgeon across 24 teams. Team formations were randomized. Each team dedicated an average of 3.5 hours within a single day to participate in a high-fidelity simulated OR environment. In the simulated OR, they encountered a series of crisis scenarios designed to test their adherence to critical and evidence-based practices. Recruitment of staff members was facilitated through sign-up sheets and random selection from those scheduled to work on designated study dates. Hospital departments arranged for staff to attend the simulation sessions instead of their regular workday. Hospital or department rules required that all anesthesia staff taking part had to have up-to-date certification



in advanced cardiac life support. Each participant only took part in one study session.

### Ethical Considerations

Ethical approval for this study was obtained from the Ministry of Health, Kuwait (IRB#: SKU-219328). Informed consent was obtained from all participants prior to their involvement in the study. Participants were informed about the study's objectives, procedures, and their rights, including the ability to withdraw at any point without any repercussions. All data collected during the study were deidentified and stored securely to ensure participant confidentiality. Data were anonymized during analysis to protect privacy, and access was restricted to authorized personnel only. No monetary or nonmonetary compensation was provided to participants for their involvement in this study. Identifiable features of participants were not captured in any images or supplementary materials.

### Primary Outcome: FTA rate

The primary outcome was the FTA rate for 47 key lifesaving processes outlined in [Multimedia Appendix 1](#). Adherence was evaluated and scored as either yes or no by 2 physician reviewers from our team (AA and RG) who observed and scored recorded simulation sessions. These sessions were recorded as synchronized videos on 2 screens for a comprehensive review. To ensure the accuracy of adherence scoring, interrater reliability was assessed. Any disagreements or uncertainties in scoring were reviewed by third reviewers (CP, HS) and were resolved. The primary variables included the checklist group and the medical crisis scenario. The primary aspect of the study was the measured FTA rates.

### Secondary Outcomes

#### Cognitive Load

We used a Polar chest strap to collect interbeat interval data from participants during scenarios with an accuracy of 1 millisecond. Previous studies have shown that a low-frequency to high-frequency (LF/HF) ratio extracted from heart rate variability is a validated proxy for cognitive load [43-45], particularly when collected using chest wraps [46]. We used NeuroKit2, a toolbox for neurophysiological signal processing [47], to extract the LF/HF ratio from data aggregated into a 1-minute time window.

### Participant Satisfaction and Usability

To evaluate the ease of use and the perceived effectiveness of the AR and paper checklists, we administered a structured survey adopted from Arriaga et al [14]. The survey assessed participants' preparedness, ease of use, readability, willingness to use the checklist in real scenarios, and perceived impact on the clinical flow during emergencies. Responses were captured on a Likert scale ranging from 1 (strongly disagree) to 5 (strongly agree), providing insights into participants' attitudes and perceptions across various aspects of checklist usage.

### Statistical Analysis

Participant characteristics were presented by descriptive statistical analysis, which reported the number and percentage of participants across different roles and years of experience. To assess the consistency in observational scoring, the agreement between two reviewers on the adherence scores was quantified using a Cohen  $\kappa$ . The Shapiro-Wilk test was used to evaluate the normality of the data distribution. ANOVA was used to compare the efficacy of interventions across 3 groups and post hoc analyses were conducted to examine the checklist's efficacy across various scenarios. Participant satisfaction and usability were analyzed using descriptive statistics and reporting means and SD. The statistical analyses were performed using SAS with all  $P$  values being 2-sided and a threshold for statistical significance set at  $P < .05$ .

## Results

### Participants

A total of 50 participants, forming 24 teams, took part in this study, which included anesthesiologists ( $n=14$ ), anesthesia residents ( $n=10$ ), OR nurses ( $n=24$ ), a surgical resident ( $n=1$ ), and a scrub nurse ( $n=1$ ). All anesthesia residents were in the early stages of their careers with 0 - 2 years of experience, and OR nurses included a more diverse range of experience, spanning from 0 - 8 years. Each team contained 1 mock surgeon and 1 surgical assistant (scrub nurse), who attended as stand-in participants to the operative field without participating in decision-making or survey completion; these stand-in staff members were not counted as participants. Participants' years of experience are summarized in [Table 1](#).


**Table 1.** Participant's role and their years of experience.

Role	Years of experience in specialty, n (%)			
	0 - 2	2 - 8	>8	Unknown
Anesthesiologist				
Attending physician ( $n=14$ )	0 (0)	7 (50)	7 (50)	0 (0)
Anesthesia resident ( $n=10$ )	10 (100)	0 (0)	0 (0)	0 (0)
Operating room nurse ( $n=24$ )	6 (25)	12 (50)	3 (12.5)	3 (12.5)
Surgical resident ( $n=1$ )	(1) 100	0 (0)	0 (0)	0 (0)
Scrub nurse ( $n=1$ )	0 (0)	1 (100)	0 (0)	0 (0)

## Adherence Rating

The assessment of adherence to key processes during the simulated scenarios demonstrated high interrater reliability among independent reviewer pairs, with Cohen  $\kappa$  values of  $\geq 0.83$  across all pairs. In instances where initial disagreement or uncertainty arose among the physician reviewers, consensus was reached through expert review with video replay. Out of a total of 595 key processes, evaluated across 24 teams for 25 key processes (excluding 8 key processes from one team that did not initiate the unexplained hypotension/hypoxia followed by an unstable bradycardia scenario), only 23 instances necessitated this expert review. The process of video replay facilitated immediate full agreement among all reviewers, highlighting the effectiveness of this approach in resolving ambiguities and ensuring accurate adherence assessment.

Figure 2. Failure to adhere to critical steps by condition type.



## Comparing Groups for Individual Crisis Scenarios

Adherence to critical steps across various scenarios demonstrated significant differences among groups, with an ANOVA test showing distinct results for asystolic cardiac arrest ( $F_{2,46}=25.07$ ;  $P<.001$ ), air embolism ( $F_{2,46}=14.90$ ;  $P<.001$ ), malignant hyperthermia ( $F_{2,46}=12.33$ ;  $P<.001$ ), and unexplained hypotension/hypoxia ( $F_{2,46}=38.39$ ;  $P<.001$ ). Post hoc analyses indicated that, across these scenarios, the AR checklist group consistently exhibited significantly lower FTA rates compared to the no checklist group, with notable differences in asystolic cardiac arrest ( $t_{23}=-6.47$ ;  $P<.001$ ), air embolism ( $t_{23}=-4.45$ ;  $P<.001$ ),

malignant hyperthermia ( $t_{23}=-4.79$ ;  $P<.001$ ), and unexplained hypotension/hypoxia ( $t_{23}=-10.57$ ;  $P<.001$ ). Comparisons between the AR and paper checklist groups were only significant for some scenarios, with slightly lower FTA rates for critical steps using the AR checklist in asystolic cardiac arrest ( $t_{23}=-2.65$ ;  $P=.014$ ) and unexplained hypotension/hypoxia ( $t_{23}=-2.10$ ;  $P=.046$ ). The paper checklist group also demonstrated significantly improved adherence over the no checklist condition in scenarios such as an air embolism ( $t_{23}=3.72$ ;  $P<.001$ ) and unexplained hypotension/hypoxia ( $t_{23}=5.40$ ;  $P<.001$ ; Figure 3).

## Comparing Groups Across All 4 Crisis Scenarios


ANOVA analysis showed significant differences in the FTA rate for critical steps among the 3 checklist groups ( $F_{2,46}=48.3$ ;  $P<.001$ ). Subsequent post hoc analysis showed the AR checklist group's mean FTA rate of 15.1% (SD 5.77%, 95% CI 13.50-16.70) was significantly lower than the paper checklist group's FTA rate of 18.32% (SD 5.65, 95% CI 16.75-19.89) and the no checklist group's FTA rate of 29.81% (SD 5.59, 95% CI 28.26-31.36). The AR group's FTA rate was significantly less than the no checklist group ( $t_{23}=-10.9$ ;  $P<.001$ ) and the paper checklist group ( $t_{23}=-2.08$ ;  $P=.048$ ). Moreover, the paper checklist group also had a significantly lower FTA rate compared to the no checklist group ( $t_{23}=-6.37$ ;  $P<.001$ ; Figure 2).

While the AR checklist group demonstrated statistically significant differences in FTA rates compared to the paper

checklist group, it is important to note that this significance was observed by a narrow margin. Given the sample size, there remains the possibility that this effect could be influenced by

chance, and further studies with larger sample sizes are necessary to confirm these findings.

**Figure 3.** Failure to adhere to critical steps by scenario and group type. AR: augmented reality.




**Cognitive Workload**

For anesthesiologists, ANOVA results showed a significant effect of the checklist type on the LF/HF ratio ( $F_{2,46}=4.88$ ;  $P=.02$ ). In pairwise comparisons, the AR checklist group had a significantly higher LF/HF ratio compared to both the paper checklist and no checklist groups, suggesting a potential increase in cognitive load when using the AR checklist ( $P<.05$ ; Figure

4). There was no significant difference in LF/HF ratio when comparing the paper checklist with no checklist groups, after adjusting for multiple comparisons. For nurses, the differences were significantly different ( $F_{2,46}=43.25$ ;  $P<.001$ ). The no checklist group had a significantly higher LF/HF ratio than the other two groups ( $P<.05$ ). The AR checklist and paper checklist groups did not differ significantly.

**Figure 4.** Low-frequency to high-frequency ratio across operating room staff roles by checklist group. AR: augmented reality; LF/HF: low frequency to high frequency; OR: operating room.



## Survey

Survey responses showed that both AR and paper checklist groups viewed their respective checklists positively (Table 2). Participants in the AR checklist group rated the checklist's ability to help them feel prepared during the emergency scenario at a mean Likert score of 4.5 (SD 0.75), and the paper checklist group rated this at 4.3 (SD 0.82), indicating no significant

difference between the groups. Participants expressed a strong willingness to use the checklists in real-life situations, with the AR group scoring a 4.6 (SD 0.70) and the paper group scoring a 4.4 (SD 0.75). When considering the disruption to the clinical flow of the operative emergency, the AR checklist group reported less disruption with a mean score of 4.5 (SD 0.90) compared to the paper checklist group's score of 4.2 (SD 1.00).

**Table .** Questionnaire response data from participants on checklist usability.

Statement	AR <sup>a</sup> checklist group (n=48), mean (SD)	Paper checklist group (n=48), mean (SD)	P value
The checklist helped me feel better prepared during the emergency scenario.	4.5 (0.75)	4.3 (0.82)	.13
The checklist was easy to use.	4.4 (0.80)	4.2 (0.85)	.09
I would use this checklist if I were presented with this operative emergency in real life.	4.6 (0.70)	4.4 (0.75)	.03
The checklist did not disrupt the clinical flow of the operative emergency.	4.5 (0.90)	4.2 (1.00)	.04
If I were having an operation and experienced this intraoperative emergency, I would want the checklist to be used.	4.7 (0.55)	4.6 (0.60)	.18

<sup>a</sup>AR: augmented reality.

## Discussion

### Principal Findings

Our findings show that AR checklist groups had a superior adherence to critical steps in crises when compared to the paper checklist groups and groups who did not use any checklist. These findings highlight AR's potential to improve OR staff's adherence to predefined protocols and ultimately improve patient outcomes. This improvement suggests that sending critical and time-sensitive information to clinicians' and OR staff's field of view may help with faster and more precise decision-making in critical situations and emergencies. Considering a day-by-day improvement in technology, this will have the potential to set the ground for an extended and more effective AR checklist intervention in many other critical scenarios. This potential benefit is in line with a comparison of the AR checklist versus the traditional checklist in other health care applications [29,30]. The benefit of AR checklists, particularly in comparison with non-AR alternatives, underscores the technology's capacity to augment traditional safety measures.

It is also important to note that while the AR checklist group had a clear superiority over the no checklist group, the margin of improvement was modest when it was compared to the paper checklist group. In this comparison, the differences were not always statistically significant across different scenarios. These findings suggest that AR technology may not offer the same improvement in all clinical scenarios over the paper checklists. Considering the low sample size and extensive subgroup analysis, it is reasonable to suggest that AR's real-world application and its superiority over conventional methods

warrant further examination. We also observed variation in team performance, as highlighted in Figure 1 of [Multimedia Appendix 1](#). Some of this variation may be attributed to an order effect, where teams became more familiar with the simulation environment over time. This potential bias should be considered when interpreting the results, and future studies could include randomization or counterbalancing to mitigate this effect.

The feedback from participants indicated a high level of acceptance and perceived utility of AR checklists in crisis scenarios, pointing to the potential for AR to integrate effectively into surgical workflows. However, the nuanced performance improvements highlight the need for a tailored approach to technological integration in health care, where the specific context and user needs dictate the effectiveness of such alternatives [48-50]. The study's results align with broader trends in medical and high-risk industries, where checklists have long been recognized for their role in promoting adherence to best practices and enhancing outcomes [51-53]. Just as checklists have transformed safety protocols in aviation and nuclear power, AR checklists hold promise for surgical settings. Nonetheless, the adaptation of these tools in medicine, particularly in the high-stakes environment of the OR, requires careful consideration of design, implementation, and training to ensure they meet the unique demands of health care providers and patients.

A key consideration emerging from our research is the differential impact of AR on the cognitive load among OR staff. Anesthesiologists using the AR checklist have shown a higher LF/HF ratio, which may be associated with a higher level of cognitive load when compared to the paper and no checklist

groups. While we initially interpreted the higher LF/HF ratio in the AR checklist group as a sign of increased cognitive burden, it is also possible that this reflects heightened cognitive engagement. The AR checklist may stimulate more focused attention on the OR environment and monitoring, compared to the paper checklist, which could be perceived as more distracting. This alternative interpretation suggests that the AR condition may enhance attentional focus in a high-stakes environment, and further research is needed to clarify the relationship between LF/HF ratio and cognitive engagement.

It is an important finding that AR technology may improve adherence but simultaneously may add a cognitive burden [54,55] that adversely affects clinicians' behavior under cognitively demanding conditions. This variability in cognitive impact across different OR roles underscores the importance of designing AR applications that are tailored to the diverse needs and cognitive capacities of surgical teams. Future studies should also include qualitative methods to capture participants' experiences with AR and paper checklists. Combining this with quantitative data will provide a more complete understanding [56].

### Limitations

This study has several limitations that should be considered. First, the study was conducted in a simulation setting that may not necessarily reflect the complexity of the OR environment. Second, our sample size was relatively small with a limited statistical power that prevented us from confidently performing

subcategory analysis and extracting minor differences between groups. Larger studies with more diverse groups of clinicians and more scenario variability are needed to allow for subgroup analyses and to look for potential impacts on certain groups of clinicians or crisis scenarios. Third, the integration of AR technology into clinical practice raises questions about cost, accessibility, and the need for specialized training [57]. The development of best practices for the implementation and customization of AR checklists will be crucial to their successful adoption in surgical care. Last, we recognize that *P* values alone should not be taken as conclusive evidence of AR's superiority. The narrow statistical margin highlights the need for further validation through larger studies to confirm its efficacy.

### Conclusion

Our study showed that the use of AR-enhanced checklists significantly improved adherence to critical procedural steps during simulated OR crises compared to both traditional paper checklists and scenarios without a checklist. These findings are promising as they may contribute to the patient's safety and outcomes. However, while the benefits of AR are promising, our findings also indicate a potential increase in cognitive load among clinicians, particularly anesthesiologists. Future studies should aim to optimize AR interfaces to minimize cognitive demands and validate these results in real-world settings. Addressing the balance between improved protocol adherence and cognitive load will be crucial for integrating AR effectively in high-stakes environments like the OR.

### Conflicts of Interest

AG is the Medical Director of Ultrasight.

### Multimedia Appendix 1

Supplementary materials on the development and application of augmented reality checklists for crisis management in clinical settings.

[DOCX File, 106 KB - [xr\\_v2i1e60792\\_app1.docx](#) ]

### References

1. Holzman RS, Cooper JB, Gaba DM, Philip JH, Small SD, Feinstein D. Anesthesia crisis resource management: real-life simulation training in operating room crises. *J Clin Anesth* 1995 Dec;7(8):675-687. [doi: [10.1016/0952-8180\(95\)00146-8](#)] [Medline: [8747567](#)]
2. Hepner DL, Arriaga AF, Cooper JB, et al. Operating room crisis checklists and emergency manuals. *Anesthesiology* 2017 Aug;127(2):384-392. [doi: [10.1097/ALN.0000000000001731](#)] [Medline: [28604405](#)]
3. Patel B, Thind A. Usability of mobile health apps for postoperative care: systematic review. *JMIR Perioper Med* 2020 Jul 20;3(2):e19099. [doi: [10.2196/19099](#)] [Medline: [33393925](#)]
4. Moorthy K, Munz Y, Forrest D, et al. Surgical crisis management skills training and assessment: a simulation[corrected]-based approach to enhancing operating room performance. *Ann Surg* 2006 Jul;244(1):139-147. [doi: [10.1097/01.sla.0000217618.30744.61](#)] [Medline: [16794399](#)]
5. Agarwala AV, Nurudeen S, Haynes AB. Perioperative checklists and handoffs. *Adv Anesth* 2015;33(1):191-217. [doi: [10.1016/j.aan.2015.07.011](#)]
6. Harari RE, Dias RD, Kennedy-Metz LR, et al. Deep learning analysis of surgical video recordings to assess nontechnical skills. *JAMA Netw Open* 2024 Jul 1;7(7):e2422520. [doi: [10.1001/jamanetworkopen.2024.22520](#)] [Medline: [39083274](#)]
7. Dias RD, Harari RE, Zenati MA, et al. A clinician-centered explainable artificial intelligence framework for decision support in the operating theatre. *Hamlyn Symp Med Robot* 2024 Jun;16:35-36. [Medline: [39076411](#)]
8. Harari R, Dias RD, Salas E, Unhelkar V, Chaspari T, Zenati M. Misalignment of cognitive processes within cardiac surgery teams. *Hamlyn Symp Med Robot* 2024 Jun;16:33-34. [Medline: [39081305](#)]



9. Lin CC, Chen YP, Chiang CC, Chang MC, Lee OKS. Real-time streaming of surgery performance and intraoperative imaging data in the hybrid operating room: development and usability study. *JMIR Med Inform* 2020 Apr 22;8(4):e18094. [doi: [10.2196/18094](https://doi.org/10.2196/18094)] [Medline: [32209528](https://pubmed.ncbi.nlm.nih.gov/32209528/)]
10. Vogt L, Klasen M, Rossaint R, Goeretz U, Ebus P, Sopka S. Virtual reality tour to reduce perioperative anxiety in an operating setting before anesthesia: randomized clinical trial. *J Med Internet Res* 2021 Sep 1;23(9):e28018. [doi: [10.2196/28018](https://doi.org/10.2196/28018)] [Medline: [34252034](https://pubmed.ncbi.nlm.nih.gov/34252034/)]
11. Harari RE, Ahmadi N, Pourfalamatoun S, Al-Taweel A, Shokoohi H. Clinician-AI collaboration for decision support in telemedicine: a randomized controlled trial study. 2024 Presented at: Proceedings of Conference on Cognitive and Computational Aspects of Situation Management 2023; Oct 16-19, 2023; Philadelphia, PA p. 81-89. [doi: [10.29007/9qxd](https://doi.org/10.29007/9qxd)]
12. Alfred MC, Cohen TN, Cohen KA, et al. Using flow disruptions to examine system safety in robotic-assisted surgery: protocol for a stepped wedge crossover design. *JMIR Res Protoc* 2021 Feb 9;10(2):e25284. [doi: [10.2196/25284](https://doi.org/10.2196/25284)] [Medline: [33560239](https://pubmed.ncbi.nlm.nih.gov/33560239/)]
13. Haynes AB, Weiser TG, Berry WR, et al. A surgical safety checklist to reduce morbidity and mortality in a global population. *N Engl J Med* 2009 Jan 29;360(5):491-499. [doi: [10.1056/NEJMsa0810119](https://doi.org/10.1056/NEJMsa0810119)] [Medline: [19144931](https://pubmed.ncbi.nlm.nih.gov/19144931/)]
14. Arriaga AF, Bader AM, Wong JM, et al. Simulation-based trial of surgical-crisis checklists. *N Engl J Med* 2013 Jan 17;368(3):246-253. [doi: [10.1056/NEJMsa1204720](https://doi.org/10.1056/NEJMsa1204720)] [Medline: [23323901](https://pubmed.ncbi.nlm.nih.gov/23323901/)]
15. Birkmeyer JD, Finks JF, O'Reilly A, et al. Surgical skill and complication rates after bariatric surgery. *N Engl J Med* 2013 Oct 10;369(15):1434-1442. [doi: [10.1056/NEJMsa1300625](https://doi.org/10.1056/NEJMsa1300625)] [Medline: [24106936](https://pubmed.ncbi.nlm.nih.gov/24106936/)]
16. McMullan RD, Urwin R, Gates P, Sunderland N, Westbrook JI. Are operating room distractions, interruptions and disruptions associated with performance and patient safety? A systematic review and meta-analysis. *Int J Qual Health Care* 2021 Apr 28;33(2):mzab068. [doi: [10.1093/intqhc/mzab068](https://doi.org/10.1093/intqhc/mzab068)] [Medline: [33856028](https://pubmed.ncbi.nlm.nih.gov/33856028/)]
17. Babayan RK. Re: crisis checklists for the operating room: development and pilot testing. *J Urol* 2012 Feb;187(2):649. [doi: [10.1016/j.juro.2011.10.071](https://doi.org/10.1016/j.juro.2011.10.071)] [Medline: [22237372](https://pubmed.ncbi.nlm.nih.gov/22237372/)]
18. Saxena S, Krombach JW, Nahrwold DA, Pirracchio R. Anaesthesia-specific checklists: a systematic review of impact. *Anaesth Crit Care Pain Med* 2020 Feb;39(1):65-73. [doi: [10.1016/j.accpm.2019.07.011](https://doi.org/10.1016/j.accpm.2019.07.011)] [Medline: [31374366](https://pubmed.ncbi.nlm.nih.gov/31374366/)]
19. Simmons WR, Huang J. Operating room emergency manuals improve patient safety: a systemic review. *Cureus* 2019 Jun 12;11(6):e4888. [doi: [10.7759/cureus.4888](https://doi.org/10.7759/cureus.4888)] [Medline: [31423368](https://pubmed.ncbi.nlm.nih.gov/31423368/)]
20. Gaba DM, Fish KJ, Howard SK, Burden A. Crisis Management in Anesthesiology: Elsevier Health Sciences; 2014:434.
21. Dryver E, Lundager Forberg J, Hård Af Segerstad C, Dupont WD, Bergenfelz A, Ekelund U. Medical crisis checklists in the emergency department: a simulation-based multi-institutional randomised controlled trial. *BMJ Qual Saf* 2021 Sep;30(9):697-705. [doi: [10.1136/bmjqs-2020-012740](https://doi.org/10.1136/bmjqs-2020-012740)] [Medline: [33597283](https://pubmed.ncbi.nlm.nih.gov/33597283/)]
22. Weinger MB. Time out! Rethinking surgical safety: more than just a checklist. *BMJ Qual Saf* 2021 Aug;30(8):613-617. [doi: [10.1136/bmjqs-2020-012600](https://doi.org/10.1136/bmjqs-2020-012600)] [Medline: [33758034](https://pubmed.ncbi.nlm.nih.gov/33758034/)]
23. Urban D, Burian BK, Patel K, et al. Surgical teams' attitudes about surgical safety and the surgical safety checklist at 10 years. *Ann Surg Open* 2021 Sep;2(3):e075. [doi: [10.1097/AS9.0000000000000075](https://doi.org/10.1097/AS9.0000000000000075)]
24. Neuhaus C, Spies A, Wilk H, Weigand MA, Lichtenstern C. "Attention everyone, time out!": safety attitudes and checklist practices in anesthesiology in Germany. A cross-sectional study. *J Patient Saf* 2021 Sep 1;17(6):467-471. [doi: [10.1097/PTS.0000000000000386](https://doi.org/10.1097/PTS.0000000000000386)] [Medline: [28574957](https://pubmed.ncbi.nlm.nih.gov/28574957/)]
25. Ebnali M, Fathi R, Lamb R, Pourfalamatoun S, Motamedi S. Using augmented holographic UIs to communicate automation reliability in partially automated driving. 2020 Presented at: Automation Experience across Domains at CHI 2020; Apr 26-27, 2020 URL: <https://ceur-ws.org/Vol-2700/paper5.pdf> [accessed 2024-11-27]
26. Harari R, Al-Taweel A, Ahram T, Shokoohi H. Explainable AI and augmented reality in transesophageal echocardiography (TEE) imaging. Presented at: 2024 IEEE International Conference on Artificial Intelligence and eXtended and Virtual Reality (AIxVR); Jan 17-19, 2024; Los Angeles, CA p. 306-309. [doi: [10.1109/AIxVR59861.2024.00051](https://doi.org/10.1109/AIxVR59861.2024.00051)]
27. Devoto L, Muscroft S, Chand M. Highly accurate, patient-specific, 3-dimensional mixed-reality model creation for surgical training and decision-making. *JAMA Surg* 2019 Oct 1;154(10):968-969. [doi: [10.1001/jamasurg.2019.2546](https://doi.org/10.1001/jamasurg.2019.2546)] [Medline: [31433467](https://pubmed.ncbi.nlm.nih.gov/31433467/)]
28. Worlikar H, Coleman S, Kelly J, et al. Mixed reality platforms in telehealth delivery: scoping review. *JMIR Biomed Eng* 2023 Mar 24;8:e42709. [doi: [10.2196/42709](https://doi.org/10.2196/42709)] [Medline: [38875694](https://pubmed.ncbi.nlm.nih.gov/38875694/)]
29. Alismail A, Thomas J, Daher NS, et al. Augmented reality glasses improve adherence to evidence-based intubation practice. *Adv Med Educ Pract* 2019;10:279-286. [doi: [10.2147/AMEP.S201640](https://doi.org/10.2147/AMEP.S201640)] [Medline: [31191075](https://pubmed.ncbi.nlm.nih.gov/31191075/)]
30. Siebert JN, Ehrler F, Gervais A, et al. Adherence to AHA guidelines when adapted for augmented reality glasses for assisted pediatric cardiopulmonary resuscitation: a randomized controlled trial. *J Med Internet Res* 2017 May 29;19(5):e183. [doi: [10.2196/jmir.7379](https://doi.org/10.2196/jmir.7379)] [Medline: [28554878](https://pubmed.ncbi.nlm.nih.gov/28554878/)]
31. Tsang KD, Ottow MK, van Heijst AFJ, Antonius TAJ. Electronic decision support in the delivery room using augmented reality to improve newborn life support guideline adherence: a randomized controlled pilot study. *Simul Healthc* 2022 Oct 1;17(5):293-298. [doi: [10.1097/SIH.0000000000000631](https://doi.org/10.1097/SIH.0000000000000631)] [Medline: [35102128](https://pubmed.ncbi.nlm.nih.gov/35102128/)]
32. Eves J, Sudarsanam A, Shalhoub J, Amiras D. Augmented reality in vascular and endovascular surgery: scoping review. *JMIR Serious Games* 2022 Sep 23;10(3):e34501. [doi: [10.2196/34501](https://doi.org/10.2196/34501)] [Medline: [36149736](https://pubmed.ncbi.nlm.nih.gov/36149736/)]



33. Puladi B, Ooms M, Bellgardt M, et al. Augmented reality-based surgery on the human cadaver using a new generation of optical head-mounted displays: development and feasibility study. *JMIR Serious Games* 2022 Apr 25;10(2):e34781. [doi: [10.2196/34781](https://doi.org/10.2196/34781)] [Medline: [35468090](https://pubmed.ncbi.nlm.nih.gov/35468090/)]
34. Kuyt K, Park SH, Chang TP, Jung T, MacKinnon R. The use of virtual reality and augmented reality to enhance cardio-pulmonary resuscitation: a scoping review. *Adv Simul (Lond)* 2021 Apr 12;6(1):11. [doi: [10.1186/s41077-021-00158-0](https://doi.org/10.1186/s41077-021-00158-0)] [Medline: [33845911](https://pubmed.ncbi.nlm.nih.gov/33845911/)]
35. Li C, Zheng B, Yu Q, Yang B, Liang C, Liu Y. Augmented reality and 3-dimensional printing technologies for guiding complex thoracoscopic surgery. *Ann Thorac Surg* 2021 Nov;112(5):1624-1631. [doi: [10.1016/j.athoracsur.2020.10.037](https://doi.org/10.1016/j.athoracsur.2020.10.037)] [Medline: [33275930](https://pubmed.ncbi.nlm.nih.gov/33275930/)]
36. Lawson J, Martin G, Guha P, et al. Effect of mixed reality on delivery of emergency medical care in a simulated environment: a pilot randomized crossover trial. *JAMA Netw Open* 2023 Aug 1;6(8):e2330338. [doi: [10.1001/jamanetworkopen.2023.30338](https://doi.org/10.1001/jamanetworkopen.2023.30338)] [Medline: [37639272](https://pubmed.ncbi.nlm.nih.gov/37639272/)]
37. Nickel F, Cizmic A, Chand M. Telestration and augmented reality in minimally invasive surgery: an invaluable tool in the age of COVID-19 for remote proctoring and telementoring. *JAMA Surg* 2022 Feb 1;157(2):169-170. [doi: [10.1001/jamasurg.2021.3604](https://doi.org/10.1001/jamasurg.2021.3604)] [Medline: [34705030](https://pubmed.ncbi.nlm.nih.gov/34705030/)]
38. Wellens LM, Meulstee J, van de Ven CP, et al. Comparison of 3-dimensional and augmented reality kidney models with conventional imaging data in the preoperative assessment of children with Wilms tumors. *JAMA Netw Open* 2019 Apr 5;2(4):e192633. [doi: [10.1001/jamanetworkopen.2019.2633](https://doi.org/10.1001/jamanetworkopen.2019.2633)] [Medline: [31002326](https://pubmed.ncbi.nlm.nih.gov/31002326/)]
39. Ebnali M, Ahmadi N, Nabiyouni E, Karimi H. AI-powered human digital twins in virtual therapeutic sessions. *Proc Int Symp Hum Factors Ergon Health Care* 2023 Mar;12(1):1-4. [doi: [10.1177/2327857923121000](https://doi.org/10.1177/2327857923121000)]
40. Ebnali M, Paladugu P, Miccile C, et al. Extended reality applications for space health. *Aerosp Med Hum Perform* 2023 Mar 1;94(3):122-130. [doi: [10.3357/AMHP.6131.2023](https://doi.org/10.3357/AMHP.6131.2023)] [Medline: [36829279](https://pubmed.ncbi.nlm.nih.gov/36829279/)]
41. Burian BK, Ebnali M, Robertson JM, et al. Using extended reality (XR) for medical training and real-time clinical support during deep space missions. *Appl Ergon* 2023 Jan;106:103902. [doi: [10.1016/j.apergo.2022.103902](https://doi.org/10.1016/j.apergo.2022.103902)] [Medline: [36162274](https://pubmed.ncbi.nlm.nih.gov/36162274/)]
42. Ziewacz JE, Arriaga AF, Bader AM, et al. Crisis checklists for the operating room: development and pilot testing. *J Am Coll Surg* 2011 Aug;213(2):212-217. [doi: [10.1016/j.jamcollsurg.2011.04.031](https://doi.org/10.1016/j.jamcollsurg.2011.04.031)] [Medline: [21658974](https://pubmed.ncbi.nlm.nih.gov/21658974/)]
43. Mizuno K, Tanaka M, Yamaguti K, Kajimoto O, Kuratsune H, Watanabe Y. Mental fatigue caused by prolonged cognitive load associated with sympathetic hyperactivity. *Behav Brain Funct* 2011 May 23;7:17. [doi: [10.1186/1744-9081-7-17](https://doi.org/10.1186/1744-9081-7-17)] [Medline: [21605411](https://pubmed.ncbi.nlm.nih.gov/21605411/)]
44. Gabaude C, Baracat B, Jallais C, Bonniaud M, Fort A. Cognitive load measurement while driving. 2012 Presented at: Human Factors and Ergonomics Society Europe Chapter Annual Meeting; Oct 10-12, 2012; Toulouse, France p. 67-80 URL: <https://hal.science/hal-01027475> [accessed 2024-12-30]
45. von Rosenberg W, Chanwimalueang T, Adjei T, Jaffer U, Goverdovsky V, Mandic DP. Resolving ambiguities in the LF/HF Ratio: LF-HF scatter plots for the categorization of mental and physical stress from HRV. *Front Physiol* 2017;8:360. [doi: [10.3389/fphys.2017.00360](https://doi.org/10.3389/fphys.2017.00360)] [Medline: [28659811](https://pubmed.ncbi.nlm.nih.gov/28659811/)]
46. Villani V, Righi M, Sabattini L, Secchi C. Wearable devices for the assessment of cognitive effort for human-robot interaction. *IEEE Sens J* 2020 Nov 1;20(21):13047-13056. [doi: [10.1109/JSEN.2020.3001635](https://doi.org/10.1109/JSEN.2020.3001635)]
47. Makowski D, Pham T, Lau ZJ, et al. NeuroKit2: a Python toolbox for neurophysiological signal processing. *Behav Res Methods* 2021 Aug;53(4):1689-1696. [doi: [10.3758/s13428-020-01516-y](https://doi.org/10.3758/s13428-020-01516-y)] [Medline: [33528817](https://pubmed.ncbi.nlm.nih.gov/33528817/)]
48. McCloskey K, Turlip R, Ahmad HS, Ghenbot YG, Chauhan D, Yoon JW. Virtual and augmented reality in spine surgery: a systematic review. *World Neurosurg* 2023 May;173:96-107. [doi: [10.1016/j.wneu.2023.02.068](https://doi.org/10.1016/j.wneu.2023.02.068)] [Medline: [36812986](https://pubmed.ncbi.nlm.nih.gov/36812986/)]
49. Rizzo MG Jr, Costello JP II, Luxenburg D, Cohen JL, Alberti N, Kaplan LD. Augmented reality for perioperative anxiety in patients undergoing surgery. *JAMA Netw Open* 2023 Aug 1;6(8):e2329310. [doi: [10.1001/jamanetworkopen.2023.29310](https://doi.org/10.1001/jamanetworkopen.2023.29310)] [Medline: [37589975](https://pubmed.ncbi.nlm.nih.gov/37589975/)]
50. Rosvall BR, Choby GW. Navigating the nose-the future of endoscopic endonasal surgical navigation. *JAMA Otolaryngol Head Neck Surg* 2024 Apr 1;150(4):326-327. [doi: [10.1001/jamaoto.2024.0025](https://doi.org/10.1001/jamaoto.2024.0025)] [Medline: [38451546](https://pubmed.ncbi.nlm.nih.gov/38451546/)]
51. Ko HCH, Turner TJ, Finnigan MA. Systematic review of safety checklists for use by medical care teams in acute hospital settings--limited evidence of effectiveness. *BMC Health Serv Res* 2011 Sep 2;11. [doi: [10.1186/1472-6963-11-211](https://doi.org/10.1186/1472-6963-11-211)] [Medline: [21884618](https://pubmed.ncbi.nlm.nih.gov/21884618/)]
52. Storesund A, Haugen AS, Flaatten H, et al. Clinical efficacy of combined surgical patient safety system and the World Health Organization's checklists in surgery: a nonrandomized clinical trial. *JAMA Surg* 2020 Jul 1;155(7):562-570. [doi: [10.1001/jamasurg.2020.0989](https://doi.org/10.1001/jamasurg.2020.0989)] [Medline: [32401293](https://pubmed.ncbi.nlm.nih.gov/32401293/)]
53. Borchard A, Schwappach DLB, Barbir A, Bezzola P. A systematic review of the effectiveness, compliance, and critical factors for implementation of safety checklists in surgery. *Ann Surg* 2012 Dec;256(6):925-933. [doi: [10.1097/SLA.0b013e3182682f27](https://doi.org/10.1097/SLA.0b013e3182682f27)] [Medline: [22968074](https://pubmed.ncbi.nlm.nih.gov/22968074/)]
54. Aitamurto T, Boin JB, Chen K, Cherif A, Shridhar S. The impact of augmented reality on art engagement: liking, impression of learning, and distraction. Presented at: Virtual, Augmented and Mixed Reality: Applications in Health, Cultural Heritage, and Industry; Jul 15-20, 2018; Las Vegas, NV p. 153-171. [doi: [10.1007/978-3-319-91584-5\\_13](https://doi.org/10.1007/978-3-319-91584-5_13)]

55. Dixon BJ, Daly MJ, Chan H, Vescan AD, Witterick IJ, Irish JC. Surgeons blinded by enhanced navigation: the effect of augmented reality on attention. *Surg Endosc* 2013 Feb;27(2):454-461. [doi: [10.1007/s00464-012-2457-3](https://doi.org/10.1007/s00464-012-2457-3)] [Medline: [22833264](https://pubmed.ncbi.nlm.nih.gov/22833264/)]
56. Weller JM, Merry AF, Robinson BJ, Warman GR, Janssen A. The impact of trained assistance on error rates in anaesthesia: a simulation-based randomised controlled trial. *Anaesthesia* 2009 Feb;64(2):126-130. [doi: [10.1111/j.1365-2044.2008.05743.x](https://doi.org/10.1111/j.1365-2044.2008.05743.x)] [Medline: [19143687](https://pubmed.ncbi.nlm.nih.gov/19143687/)]
57. Borna A, Mousavi SZ, Fathollahzadeh F, Nazeri A, Harari RE. Applications of augmented and virtual reality in enhancing communication for individuals who are hard of hearing: a systematic review. *Am J Audiol* 2024 Oct 7:1-17. [doi: [10.1044/2024\\_AJA-24-00056](https://doi.org/10.1044/2024_AJA-24-00056)] [Medline: [39374491](https://pubmed.ncbi.nlm.nih.gov/39374491/)]

## Abbreviations

**AR:** augmented reality

**FTA:** failure to adhere

**LF/HF:** low-frequency to high-frequency

**OR:** operating room

*Edited by L Riedemann, T Leung; submitted 21.05.24; peer-reviewed by B Robinson, K Bielka; revised version received 29.10.24; accepted 29.10.24; published 06.01.25.*

*Please cite as:*

*Ebnali Harari R, Altaweel A, Anderson E, Pozner C, Grossmann R, Goldsmith A, Shokoohi H*

*Augmented Reality in Enhancing Operating Room Crisis Checklist Adherence: Randomized Comparative Efficacy Study*

*JMIR XR Spatial Comput* 2025;2:e60792

URL: <https://xr.jmir.org/2025/1/e60792>

doi: [10.2196/60792](https://doi.org/10.2196/60792)

© Rayan Ebnali Harari, Abdullah Altaweel, Erik Anderson, Charles Pozner, Rafael Grossmann, Andrew Goldsmith, Hamid Shokoohi. Originally published in *JMIR XR and Spatial Computing* (<https://xr.jmir.org>), 6.1.2025. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in *JMIR XR and Spatial Computing*, is properly cited. The complete bibliographic information, a link to the original publication on <https://xr.jmir.org/>, as well as this copyright and license information must be included.

---

Publisher:  
JMIR Publications  
130 Queens Quay East.  
Toronto, ON, M5A 3Y5  
Phone: (+1) 416-583-2040  
Email: [support@jmir.org](mailto:support@jmir.org)

---

<https://www.jmirpublications.com/>